

## Notes on Alien *Bromus* Grasses in Taiwan

Ming-Jer Jung<sup>(1)</sup>, Gwo-Ing Liao<sup>(1)</sup> and Chang-Sheng Kuoh<sup>(1,2)</sup>

(Manuscript received 15 September, 2005; accepted 1 November, 2005)

**ABSTRACT:** *Bromus carinatus* Hook. & Arn., *Bromus hordeaceus* L., *Bromus pubescens* Muhl. ex Willd. and *Bromus secalinus* L. were recently found at middle elevations of southern and central Taiwan, respectively. We present taxonomic treatments, distribution map, and line-drawings of these introduced alien brome grasses.

**KEY WORDS:** *Bromus carinatus*, *Bromus hordeaceus*, *Bromus pubescens*, *Bromus secalinus*, Flora, Gramineae, Poaceae, Taiwan.

### INTRODUCTION

In participation of a long term project, we have been collecting many plant specimens in Taiwan for drug screening since 2002. Recently, we found several species new to the Flora of Taiwan. The genus *Bromus* L. has ca. 150 species in both hemispheres (Clayton and Renvoize, 1986), and four taxa were recorded in the Flora of Taiwan, including two naturalized species (Hsu, 1978; Kuoh and Chen, 2000; Huang et al., 2003). *Bromus formosanus* Honda, which was treated as endemic species in Taiwan, occurs in Sabah and New Guinea, too (J. F. Veldkamp, personal communication). *B. morrisonensis* Honda occurring in Malesia and Luzon was also treated as endemic species in Taiwan (Veldkamp et al., 1991). Besides, *B. rigidus* Roth in Roemer and Usteri. occurring at An-ma-shan forest trail 43 K (Kuoh and Chen, 2000) must be corrected to *B. diandrus* Roth.

We treated four additional alien grasses, i.e. *Bromus carinatus* Hook. & Arn., *Bromus hordeaceus* L., *Bromus pubescens* Muhl. ex Willd. and *Bromus secalinus* L. with distributional maps (Fig. 1) and illustrations. A key to brome grasses in Taiwan is also provided here.

### TAXONOMIC TREATMENT

1. *Bromus carinatus* Hook. & Arn., Bot. Beechey Voy. 403. 1840; Hitchcock A. S., Manual of grasses of the United States, 35-36, t. 7. 1951; Osada T., Illustrated Grasses of Japan, 398-399, 1993. 長芒扁雀麥 Fig. 2

Culms erect, to 100 cm tall, 5-7-noded; nodes articulate, glabrous, exposed. Ligule triangular, ca. 3 mm long, membranous, abaxial surface pubescent; sheath glabrous to pubescent, shorter than blades; blade up to 30 × 0.5-1.5 cm, 11-veined with one main vein. Inflorescence a panicle, spreading, with ca. 4 nodes, 30-40 cm long; basal branches ca. 15 cm long. Spikelet oblong, 2-4 cm × 3-6 mm, with 2-7 florets. Lower glume lanceolate, 0.9-1.2 cm long, 3-veined, margin membranous, apex acuminate. Upper glume lanceolate, 1.2-1.4 cm long, margin membranous, 7-veined, apex acuminate or with one short awn. Lemma lanceolate, 1.5-1.8 cm long, 7-veined with keeled central vein, apex obtuse; awn arising from center of apex, 6-9 mm long. Palea lanceolate, apex acuminate, 2-keeled, keels hispid. Anthers ca. 4 mm long. Ovary ca. 1.5 mm long, pilose, hairs ca. 0.75 mm long.

Distribution and notes: *Bromus carinatus* (California brome) is distributed in western America (Hitchcock, 1951). In Taiwan, it occurs at Kuaigu, Tienchi and the entrance of the Li-guan Tunnel, along the Southern Cross-Island Highway (Taiwan Highway No. 20), and ca. 1800 m elevation (Fig. 1). It is associated with many grasses such as *Brachypodium sylvaticum* (Huds.) P. Beauv., *Bromus hordeaceus* L., *Dactylis glomerata* L., *Festuca arundinacea* Schreb., *Lolium perenne* L., *Poa annua* L., *Polypogon fugax* Nees ex Steud., and *Setaria plicata* (Lam.) T. Cooke. *Bromus carinatus* can be distinguished from *B. catharticus* by the longer awn and lesser compressed spikelet.

Specimens examined: Kaohsiung Co.: Tau-yuan Hsiang, Entrance of Li-guan Tunnel, M.-J. Jung, x061010, x061012 (NCKU); Jhongjhihguan, T.-H. Hsieh & C.-F. Lin s. n., Jun 20, 2004 (NTNTC); Tienchi, M.-J. Jung, x071103, x071114, x071115, x071116, x071117 (NCKU); Kuaigu, M.-J. Jung, x100501, x100502, x100503 (NCKU).

1. Department of Life Sciences, National Cheng-Kung University, 1, Ta-Hsueh Road, Tainan 701, Taiwan.

2. Corresponding author. Email: kuohpopo@gmail.com




Fig. 1. Distribution of *Bromus carinatus* (☆) and *B. hordeaceus* (★), *B. pubescens* (▼) and *B. secalinus* (□) in Taiwan.

2. *Bromus hordeaceus* L., *Sp. Pl.* 1: 77. 1753; Pavlick L. E., *Bromus L. of North America*, 78-81. 1995; Smith in Cafferty et al., *Taxon* 49(2): 248. 2000. 毛雀麥 Fig. 3

*Bromus mollis* L. *Sp. Pl.* (ed. 2) 1: 112. 1762; Hitchcock A. S., *Manual of grasses of the United States*, 49-50, t. 29. 1951; Osada T., *Illustrated Grasses of Japan*, 390-391, 1993

Culms annual, erect, to 30 cm tall; stem long tomentose. Leaf sheath long tomentose; ligule ca. 0.8 mm long, membranous; blade linear, ca. 5 cm long. Inflorescence spike-like with few spikelets; or a panicle. Lower glume elliptical, ca. 6 mm long, tomentose, 7-veined. Upper glume ovate, ca. 7 mm long, tomentose, 9-veined. Lemma obovate, ca. 8 mm long, 7-veined, apex bifid, tomentose, with one awn; awn arising from 1/5 of apex, 7-8 mm long. Palea oblong, ca. 6.5 mm long, 2-keeled, apex obtuse and ciliate; keels strigose and hispid. Caryopsis oblong, ca. 6 mm long, base caudate, adnate with lemma and palea when mature.

Distribution and notes: The common name of *Bromus hordeaceus* is soft chess, distributed in North America and Europe. In Taiwan, it occurs at the entrance of Li-Guan Tunnel (Fig. 1), the same site as *B. carinatus* mentioned above. There are several subspecies, our specimens belong to subsp. *hordeaceus*. (J. F. Veldkamp, personal communication).

Specimens examined: Kaohsiung Co.: Tau-yuan Hsiang. Li-Guan tunnel, *M.-J. Jung*, x061003, x061004 (NCKU).

3. *Bromus pubescens* Muhl. ex Willd., *Enum. Pl.* 120. 1809; Pavlick L. E., *Bromus L. of North America*, 58 - 59. 1995. 短毛雀麥 Fig. 4

Culms to 150 cm tall, rhizomatous; stems erect; lower sheaths pubescent, retrorsely pilose, upper sheaths glabrous; ligule ca. 2 mm long, membranous; blades up to 30 × 1.3 cm. Spikelet lanceolate, with 7-13 florets. Lower glume lanceolate, 8-9 mm long, with 3-5 veins. Upper glume lanceolate, ca. 1.1 mm long, 7-veined; rachilla pubescent. Lemma oblanceolate, 7-15 mm long; abaxial surface pubescent, margin long ciliate, ciliate hairs ca. 1 mm long, apex bifid, with one awn. Palea 2-keeled, keels shortly hispid. Anthers ca. 0.8 mm long. Ovary apically pubescent.

Distribution and notes: The common names of *Bromus pubescens* Muhl. ex Willd. include: Canada brome, Canadian brome, hairy brome, hairy woodland brome, and woodland brome. In Taiwan, it was found beneath Fu-ci trees, along the roadside of Taiwan Highway No. 21, at ca. 2400 m (Fig. 1). *Agrostis infirma* Buse, *Agrostis stolonifera* L., *Dactylis glomerata* L., *Dendranthema arisanense* (Hayata) Y. Ling & C. Shih, *Festuca arundinacea* Schreb., *Poa annua* L., and *Senecio inaequidens* DC. could be found at the same site.

Specimens examined: Nantou Co.: Shin-yi Hsiang. Fuci tree, *M.-J. Jung*, x060601, x060603, x062704, x062705 (NCKU).

4. *Bromus secalinus* L., *Sp. Pl.* 1: 76. 1753; Hitchcock A. S., *Manual of Grasses of the United States*, 47-49, t. 27. 1951; Osada T., *Illustrated Grasses of Japan*, 388-389, 1993. 歐雀麥 Fig. 5




Fig. 2. *Bromus carinatus* Hook. & Arn. A: Habit. B: Ligule. C: Floret with rachilla. D: Lower glume. E: Upper glume. F: Lemma. G: Palea. H: Anther. I: Pistil.




Fig. 3. *Bromus hordeaceus* L. A: Habit. B: Ligule. C: Adaxial view of floret. D: Lower glume. E: Upper glume. F: Lemma. G: Palea. H: Anther. I: Caryopsis.




Fig. 4. *Bromus pubescens* Muhl. ex Willd. A: Habit. B: Ligule. C: Nodes of inflorescence. D: Lower glume. E: Upper glume. F: Adaxial view of floret. G: Lemma. H: Palea. I: Anther. J: Caryopsis.




Fig. 5. *Bromus secalinus* L. A: Habit. B: Ligule. C: Blade apex. D: Adaxial view of floret. E: Lower glume. F: Upper glume. G: Lemma. H: Palea. I: Lodicules. J: Anther. K: Pistil.

Culms erect, to 30 cm tall. Sheath pubescent. Ligule ca. 1 mm long, membranous, apex irregularly dentate. Blade up to 20 × 1 cm, pubescent. Panicle simple to spreading, ca. 15 cm long; lower branches declinate. Spikelet ca. 3 cm long, with 7-9 florets; lower glume lanceolate, ca. 5.5 mm long, glabrous, 5-veined. Upper glume lanceolate, ca. 7 mm long, glabrous, 7-veined. Lemma obovate, ca. 9 mm long, glabrous, 7-veined, apex bifid, with one awn; awn arising from submargin of lemma apex, 8-10 mm long. Palea ca. 8 mm long, membranous, with 2-keeled, long hispid on keels. Anthers 3, ca. 1 mm long. Ovary pubescent, apex hispid.

Distribution and notes: The common names of *Bromus secalinus* are cheat, chess, and rye brome. It was introduced to America from Europe (Hitchcock, 1951). In Taiwan, it occurs near the Lu-lin sacred tree, along the Taiwan Highway No. 18, at ca. 2400 m (Fig. 1). It grows together with many grasses such as *Dactylis glomerata* L., *Festuca arundinacea* Schreb., *Miscanthus sinensis* Anders., *Poa annua* L., and *Vulpia myuros* (L.) Gmel.

Specimens examined: Nantou Hsien: Shin-yi Hsiang. Lulin Lodge, T.-H. Hsieh, 3030 (NTNTC); Lu-lin sacred tree, M.-J. Jung, x053005, x062709, x062710, x062711 (NCKU).

5. *Bromus diandrus* Roth, Bot. Abh. Beobacht. 44. 1787; Veldkamp J. F., Blumea 35(2): 489. 1991.

硬雀麥

*Bromus rigidus* Roth in Roemer & Usteri., Mag. Bot. 10: 21. 1790; Kuo C.-M., Taiwania 24: 22. 1979; Osada T., Ill. Grasses Jap. Enl. Ed. 382. 1993; Chen C.-H., & Kuoh C.-S., Taiwania 45: 311-322. 2000

### Key to the species

1. Spikelets compressed, glumes and lemmas keeled.
1. Spikelets terete, glumes and lemmas not keeled.
  2. Lemmas mostly 11-veined, awnless or with a short awn, awn shorter than 5 mm; paleaelliptical ..... *B. catharticus*
  2. Lemmas 5-7-veined, awned, awn longer than 5 mm; palea oblong. .... *B. carinatus*
  3. Lemma awn longer than 5 cm ..... *B. diandrus*
  3. Lemma awn shorter than 5 cm
    4. Lemmas board, elliptical, obovate to oblanceolate
    4. Lemmas narrow, lanceolate to oblong
      5. Lemmas glabrous ..... *B. secalinus*
      5. Lemmas pubescent
        6. Lemmas tomentose, awn arising from submargin of lemma apex. .... *B. hordeaceus*
        6. Lemmas margin ciliate, awn arising from lemma apex ..... *B. pubescens*
        7. Lower glumes 1-nerved; upper part of lemma glabrous, lower part of lemma loose hairy; awn nearly as long as lemma ..... *B. morrisonensis*
        7. Lower glumes 3-nerved; lemma dense hairy, awn shorter than 1/2 length of lemma ..... *B. formosanus*

### ACKNOWLEDGEMENTS

The authors thank Dr. Anthony R. Brach of the Harvard University Herbaria, Cambridge (Mass.), U.S.A., and J. F. Veldkamp of the National Herbarium Nederland, Leiden, the Netherlands, for critical reading of the manuscript. This work was supported by a grant, NSC 93-2323-B 006-004, from the National Science Council, Taiwan.

### LITERATURE CITED

- Cafferty, S., E. C. Jarvis and N. J. Turland. 2000. Typification of Linnaean plant names in the Poaceae (Gramineae). *Taxon* **49**: 239-260.
- Chen, C.-H. and C.-S. Kuoh. 2000. The Genus *Bromus* L. (Poaceae) in Taiwan: a DELTA Database for Generating Key and Descriptions. *Taiwania* **45**: 311-322.
- Clayton, W. and S. Renvoize. 1986. Genera graminarum, Grasses of the world, H. M. S. O., London. London, UK. p. 144.
- Hitchcock, A. S. 1951. Manual of grasses of the United States. ed. 2, rev. by A. Chase, U.S. Dep. Agr. Misc. Publ., USA. pp. 31-56.
- Huang, T.-C. 2003. In: Huang, T.-C. et al. (eds.), Flora of Taiwan, 2nd ed. 6: 121. Editorial Committee, Dept. of Bot., NTU, Taipei, Taiwan.
- Hsu, C.-C. 1978. Gramineae. In: Huang, T.-C. et al. (eds.), Flora of Taiwan, 1st ed. 5: 429-433. Editorial Committee, Dept. Bot., NTU, Taipei, Taiwan.
- Kuo, C.-M. 1979. Three Newly Recorded Naturalized Plants of Taiwan — *Mollugo verticillata*, *Festuca elatior* and *Bromus rigidus*. *Taiwania* **24**: 22-26.
- Kuoh, C.-S. and C.-H. Chen. 2000. Pooideae of Gramineae (Poaceae). In: Huang, T.-C. et al. (eds.), Flora of Taiwan, 2nd ed. 5: 370-374. Editorial Committee, Dept. Bot., NTU, Taipei, Taiwan.
- Osada, T. 1993. Illustrated Grasses of Japan. Heibonsia Ltd., Tokyo, Japan. pp. 376-399.
- Pavlick, L. E. 1995. *Bromus* L. of North America. Royal British Columbia Museum, Victoria, Canada. 160pp.
- Veldkamp, J. F., M. Eriks and S. S. Smit. 1991. *Bromus* (Gramineae) in Malesia. *Blumea* **35**: 483-497.

## 臺灣外來雀麥屬禾草增註

鍾明哲<sup>(1)</sup>、廖國嫻<sup>(1)</sup>、郭長生<sup>(1,2)</sup>

(收稿日期：2005 年 9 月 15 日；接受日期：2005 年 11 月 1 日)

### 摘 要

長芒扁雀麥 (新擬中名) (*Bromus carinatus* Hook. & Arn.)、毛雀麥 (*Bromus hordeaceus* L.)、短毛雀麥 (新擬中名) (*Bromus pubescens* Muhl. ex Willd.)、及歐雀麥 (新擬中名) (*Bromus secalinus* L.) 分別為近期在臺灣南部及中部中海拔山區所發現之外來雀麥屬植物。本文提供此四種植物的描述、形態及分布。

關鍵詞：長芒扁雀麥、毛雀麥、短毛雀麥、歐雀麥、禾本科、臺灣、分類。

---

1. 國立成功大學生命科學系，701 台南市東區大學路 1 號，臺灣。

2. 通信作者。Email: kuohpopo@gmail.com