

## *Austro eupatorium inulifolium* (Kunth) King & Robinson (Asteraceae), a Newly Naturalized Plant in Taiwan

Tsai-Wen Hsu<sup>(1,2)</sup>, Ching-I Peng<sup>(3)</sup> and Chiu-Mei Wang<sup>(4,5)</sup>

(Manuscript received 21 July, 2005; accepted 9 November, 2005)

**ABSTRACT:** In recent years we have witnessed the naturalization of many alien species of Asteraceae in Taiwan. In this report we document the recent naturalization of the Neotropical *Austro eupatorium inulifolium* (Kunth) R. M. King & H. Rob. at 1,300-1,400 m altitudes in the Central Mountain Range of Taiwan. We provide a description, a line drawing, and photographs to aid in identification.

**KEY WORDS:** Asteraceae, *Austro eupatorium inulifolium*, Naturalized plant, Taiwan, Taxonomy.

### INTRODUCTION

The tribe Eupatorieae (Asteraceae) consists of approximately 2,400 species in 170 genera (Bremer, 1994). Plants of this tribe are characterized by discoid heads with white, reddish or blue corollas. While they are concentrated in Central and South America, they are also represented in North America and, to a less extent, in the Old World (Bremer, 1994). A number of the species in Eupatorieae are widespread weeds, e.g., *Mikania micrantha* Kunth, a major weed in tropical Southeast and South Asia; *Ageratum conyzoides* L., a common weed of crops and gardens in the tropics worldwide; *Ageratina adenophora* (Spreng.) R. M. King & H. Rob. and *Ageratina riparia* (Regel) R. M. King & H. Rob. rampant in Asia (Peng et al., 1998b); *Praxelis clematidea* R. M. King & H. Rob. naturalized to northern Queensland, southern China and Taiwan; and *Chromolaena odorata* (L.) King & Robins in Australia, Africa, and Southeast and South Asia (McFadyen, 2003; Peng and Yang, 1998). In Taiwan, the tribe Eupatorieae is represented by three indigenous genera, *Adenostemma*, *Eupatorium* and *Mikania* and five genera adventive from the New World, i.e. *Ageratina*, *Ageratum*, *Austro eupatorium*, *Chromolaena* and *Praxelis*.

The genus *Austro eupatorium* R. M. King and H. Rob. comprises 13 species distributed in the Andean South America northward to Panama and Trinidad, with the center of diversity in eastern South America westward to Bolivia (King and Robinson, 1987). With the presence of glands and lack of setulae on achenes, *Austro eupatorium* is closest to *Eupatorium*, differing by the large carpodium with large thin-walled cells and the strongly ornamented walls of the anther collar (King and Robinson, 1987).

### TAXONOMIC TREATMENT

*Austro eupatorium inulifolium* (Kunth) R. M. King & H. Rob., Phytologia 19(7): 434. 1970, "inulaefolium". 假澤蘭(新擬) Figs. 1 & 2

*Eupatorium inulifolium* Kunth, Nova Gen. Sp. Pl. 4: 85. 1818.

Perennial, erect herbs or shrubs, to 2-3 m tall. Stems terete, often striate. Leaves simple, opposite below, becoming subopposite or alternate above; petiole 1-2 cm, puberulent; blades ovate to narrowly oblong, 7-14 cm long, 2-6 cm wide, acuminate at apex, base often decurrent onto petiole, the margins serrulate to serrate, 3-veined from well above base, the upper surface strigose, the lower surface puberulent. Inflorescence a corymbose panicle, terminal or arising from upper nodes, heads short-pedicellate. Involucral bracts 12-18, subimbricate, cylindric-obconic, in 3-4 series, persistent, 3-5-striate, elliptic to ovate. Florets 7-13 in a head, corolla white, bisexual, narrowly funnelliform with a narrow tube, ca. 4 mm long, shortly 5-lobed, the lobes triangular, glabrous or lobes sometimes glandular. Anthers included,

1. Endemic Species Research Institute, 1, Ming-Sheng E. Rd, Chi-Chi, Nantou 552, Taiwan.
2. Department of Life Sciences, Cheng-Kung University, 1, Ta-Hsueh Rd, Tainan 701, Taiwan.
3. Herbarium, Research Center for Biodiversity, Academia Sinica, 128, Sect. 2, Academia Rd, Nangang, Taipei 115, Taiwan.
4. Department of Botany, National Museum of Natural Science, 1, Kuan-Chien Rd, Taichung 404, Taiwan.
5. Corresponding author Chiu-Mei Wang, Tel: 886-4-23226940 ext. 520; Email: cmwang@mail.nmns.edu.tw




Fig. 1. *Austroeupatorium inulifolium* (Kunth) King & Robinson. 1: Habit. 2: Head. 3: Stamens. 4: Floret. 5: Achene. 6: Style branches.



Fig. 2. *Austroeupatorium inulifolium* (Kunth) King & Robinson. A: Habitat. B: Habit. C: Inflorescences.

filaments slender, collars narrowly cylindrical, the apical appendages ovate-oblong, longer than wide. Style erect and exserted, the base puberulous, branches filiform. Achenes elongate obconical, ca. 2 mm long, 5-ribbed, glanduliferous, without setulae, carpodium distinct, pappus of many slender

scabrous bristles, persistent, about as long as corolla and much longer than the achene.

Specimen examined: TAIWAN. Nantou County: Jenai Hsiang, Wushe-Yushih, near Jenai Junior High School, alt. 1,400 m, 18 Sep 2001, C. M. Wang & C. Y. Li 5270 (HAST, TNM), 5 Oct 2001, C. M. Wang & C. Y. Li 5284 (TNM); Jenai Hsiang, Lihsing Industrial Road, alt. 1,315 m, 21 Jan 2003, C. M. Wang & Y. M. Hsu 6545 (TNM).

Distribution and notes: In recent years we have witnessed the naturalization of many alien species of Asteraceae (Boufford and Peng, 1993; Peng, et al., 1998b; Peng and Yang, 1998; Peng et al., 1998a; Chen et al., 1999; Tseng and Peng, 2004) in Taiwan. *Austro eupatorium inulifolium* is widely distributed in Panama, Colombia, Venezuela, Ecuador, Peru, Bolivia, Brazil, Uruguay, to Argentina, and adventive in Indonesia, Sumatra, Ceylon (King and Robinson, 1987) and Sri Lanka (McFadyen, 2003). In Taiwan, it occurs at ca. 1,300-1,400 m altitudes in the Central Mountain Range. It can reach 2 m tall on open slopes, along road cuts or landslide areas, where it is associated with other weedy plants such as *Conyza sumatrensis* (Retz.) Walker, *Galinsoga quadriradiata* Ruiz & Pav., *Spergula arvensis* L., *Eupatorium cannabinum* L. ssp. *asiaticum* Kitamura, *Sonchus oleraceus* L., *Solanum mauritianum* Scopoli, *Rumex obtusifolius* L., *Solanum americanum* Miller, *Polygonum multiflorum* Thunb. var. *hypoleucum* (Ohwi) Liu, *Polygonum chinense* L., *Rubus parvifolius* L. Flowering period is between September and January.

*Austro eupatorium inulifolium* is an aggressive species that rapidly colonizes areas cleared for planting new crops, agricultural fields, fallow fields, waste lands and roadsides.

#### ACKNOWLEDGEMENTS

This work was supported by research grants from the Council of Agriculture, Taiwan.

#### LITERATURE CITED

- Boufford, D. E. and C.-I. Peng. 1993. *Soliva* Ruiz & Pavon (Anthemideae, Asteraceae) in Taiwan. Bot. Bull. Acad. Sin. **34**: 347-352.
- Bremer, K. 1994. Asteraceae, Cladistics & Classification. Timber Press, Portland, Oregon, USA. pp. 663.
- Chen, S.-H., M.-J. Wu and S.-M. Li. 1999. *Centratherum punctatum* Cass. ssp. *fruticosum*, a newly naturalized plant in Taiwan. Taiwania **44**: 299-305.
- King, R. M. and H. Robinson. 1987. The genera of the Eupatorieae (Asteraceae). Monographs in Systematic Botany, Missouri Botanical Garden **22**: 1-581.
- McFadyen, R. C. 2003. *Chromolaena* in Southeast Asia and the Pacific. In: da Costa, H. et al. (eds.), Agriculture: New Directions for a New Nation – East Timor (Timor-Leste). ACIAR Proceedings No. 113, pp. 130-134. Canberra, Australia.
- Peng, C.-I. and K.-C. Yang. 1998. Unwelcome naturalization of *Chromolaena odorata* (Asteraceae) in Taiwan. Taiwania **43**: 289-294.
- Peng, C.-I., C.-H. Chen, W.-P. Leu and H.-F. Yen. 1998a. *Pluchea* Cass. (Asteraceae: Inuleae) in Taiwan. Bot. Bull. Acad. Sin. **39**: 287-297.
- Peng, C.-I., K.-F. Chung and W.-P. Leu. 1998b. Notes on three newly naturalized plants (Asteraceae) in Taiwan. Taiwania **43**: 320-329.
- Tseng, Y.-H. and C.-I. Peng. 2004. *Ambrosia psilostachya* DC. (Asteraceae) a newly naturalized plant in Taiwan. Endemic Sp. Res. **6**: 49-52.

## 臺灣新歸化菊科植物—假澤蘭

許再文<sup>(1,2)</sup>、彭鏡毅<sup>(3)</sup>、王秋美<sup>(4,5)</sup>

(收稿日期：2005 年 7 月 21 日；接受日期：2005 年 11 月 9 日)

### 摘 要

近年來有許多的外來菊科植物在臺灣歸化。本研究提出一種歸化於臺灣中部山區，原產於新熱帶的菊科植物—假澤蘭 (*Austro eupatorium inulifolium* (Kunth) King & Robinson)，除了描述其分類性狀外，並提供植物繪圖與彩色照片。

關鍵詞：菊科、假澤蘭、歸化植物、臺灣、分類。

- 
1. 農委會特有生物研究保育中心，552 南投縣集集鎮民生東路 1 號，臺灣。
  2. 國立成功大學生命科學系，701 台南市大學路 1 號，臺灣。
  3. 中央研究院生物多樣性研究中心標本館，115 台北市南港區研究院路 2 段 128 號，臺灣。
  4. 自然科學博物館植物學組，404 台中市館前路 1 號，臺灣。
  5. 通信作者。Tel: 886-4-23226940 ext. 520; Email: cmwang@mail.nmns.edu.tw