

Leptolejeunea picta Herz., a Liverwort New to Taiwan

Jia-Dong Yang^(1,2) and Shan-Hsiung Lin^(2,3)

(Manuscript received 20 March 2008; accepted 2 June 2008)

ABSTRACT: *Leptolejeunea picta* Herz. is reported as a new liverwort species to Taiwan. A revised species description and illustration are provided along with information on the distribution and habitat for this species.

KEY WORDS: *Leptolejeunea picta*, Lejeuneaceae, liverwort, new record, Taiwan.

INTRODUCTION

There are six species of the genus *Leptolejeunea* in Taiwan (Lin, 2000; Piippo, 1990). The first author collected two specimens in Chikeshan, Yuli Township, Hualien County during in 2007. After checking references and specimens we determined that the Chikeshan collections represent *L. picta* Herz., a species new for the Taiwan bryoflora. This species is similar to *L. elliptica* (Lehm. & Lindenb.) Schiffn., the most common species of *Leptolejeunea* in Taiwan. Both are very small plants with leaves obliquely spreading, and leaf lobes ovate and apically obtuse; however, *L. picta* differs from *L. elliptica* in its dark-red ocelli scattered over the leaf lobes, lobules and underleaves. *Leptolejeunea picta* is a tropical species distributed from Indo-China, Sumatra, Borneo, New Guinea, and Ryukyu Archipelago (Mizutani, 1978). Voucher specimens for this study are deposited at the herbarium of Endemic Species Research Institute (TAIE).

TAXONOMIC TREATMENT

Leptolejeunea picta Herz., Flora 35: 430 (1942).

斑點薄鱗蘚 Fig. 1

Plants small, lightly green when fresh, dark brown when dried. Shoots 0.50-0.60 mm wide, irregularly pinnately branched. Stems up to 10 mm long, 35.6-52.9 μ m in diameter. Ventral merophytes of stem 2 cells wide, rhizoids numerous, fasciculate. Leaves imbricate, leaf lobe flat, ovate, 0.31-0.34 mm

long, 0.16-0.17 mm wide, apex obtuse to rounded, margin entire. Lobe cells thin-walled, trigones small, marginal cells 14.9-20.2 \times 10.4-14.4 μ m, median cells 20.1-31.6 \times 10.8-20.7 μ m, basal cells slightly larger than median ones. Cuticle smooth. Oil bodies 6-15 per median cell of leaf lobe, of the compound type. Ocelli dark red when plants dried, scattered, basal ocelli 1-2, very large, 33.8-79.9 \times 14.8-24.9 μ m, median ocelli 16.9-27.2 \times 14.1-16.9 μ m. Lobule 1/3-1/2 the length of the lobe, oblong, inflated, the first tooth unicellular, obtuse, the second tooth obsolete, the hyaline papilla on the proximal side of the first tooth. Underleaves 0.76-1.16 mm long, 0.54-1.04 mm wide, distant, transversely inserted, the basal portion usually with one blunt tooth on each lateral margin, the upper margin usually composed of 2 ocelli, the lobe linear, 2-celled.

Distribution: Indo-China, Sumatra, Borneo, New Guinea, and Ryukyu Archipelago. New to Taiwan.

Specimens examined: TAIWAN, Hualien: Yuli, Chikeshan, near the Fute Temple, creek bank, mixed hardwood forest, epiphyllous, on leaves of fern [*Osmunda banksiifolia* (Presl) Kuhn], ca. 850 m alt., E 121°23'; N 23°23', Oct. 18, 2007, J.D. Yang 4734g (TAIE), J.D. Yang 4735f (TAIE).

Comment: Growing on the leaves of *Osmunda banksiifolia*, *L. picta* usually occupied the middle part of the pinnule and associated with many common and wild distributed epiphyllous liverworts, such as *L. elliptica*, *Cololejeunea floccosa* (Lehm. & Lindenb.) Schiffn., *C. goebelli* (Gotteshe ex Schiffn.) Schiffn., and *C. appressa* (A. Evans) Benedix. This species is easily separated from other members of the genus *Leptolejeunea* by the ocelli scattered over the leaf lobes, lobules and underleaves. The ocelli turned dark red in color when plants dried, also a useful diagnostic character.

1. Endemic Species Research Institute, 1, Min-Sheng E. Rd., Chi-Chi Town, Nantou County 55244, Taiwan.

2. Department of Life Science, Tunghai University, 181, Sect. 3, Taichung-Kang Rd., Taichung 40704, Taiwan.

3. Corresponding author. Tel: 886-4-23590121 ext. 32406; Email: slin@thu.edu.tw




Fig. 1. *Leptolejeunea picta* Herz. a & c: Marginal cells of leaf lobes. b & e: Median cells with ocelli and oil bodies of leaf lobes. d: Apex cells of leaf lobe. f: Portion of sterile plant, ventral view. g & m: Underleaves. h: Leaf, ventral view. l & j: Leaf lobules. k: Leaf of cladium. l: Cladium, ventral view. All drawn from J.D. Yang 4734g.

ACKNOWLEDGEMENTS

Thanks are due to James R. Shevock for reviewing the manuscript and making useful suggestions. We are grateful to Miss Hsiu-Jane Chen and Miss Kui-Chu Chen who assisted in field collections and plate preparation. We also thank for the financial support by National Science Council (NSC 96-2621-B-329-001) and Council of Agriculture, Taiwan.

LITERATURE CITED

- Lin, S.-H. 2000. The liverwort flora of Taiwan. The Council of Agriculture, Taipei, Taiwan. (In Chinese)
- Mizutani, M. 1978. Lejeuneaceae from Ishigaki and Iriomote Islands of Ryukyu Archipelago. *Journ. Hattori Bot. Lab.* **44**: 121-136.
- Piippo, S. 1990. Annotated catalogue of Chinese Hepaticae and Anthocerotae. *Journ. Hattori Bot. Lab.* **68**: 1-192.

臺灣新紀錄蘚類—斑點薄鱗蘚

楊嘉棟^(1,2)、林善雄^(2,3)

(收稿日期：2008年3月20日；接受日期：2008年6月2日)

摘 要

本文報導新近確認的一種臺灣新紀錄蘚類—斑點薄鱗蘚(新擬)。文中附有形態描述、植物線條圖、棲地、分布及引證標本等。

關鍵詞：斑點薄鱗蘚、細鱗蘚科、蘚類、新紀錄、臺灣。

1. 行政院農業委員會特有生物研究保育中心，55244 南投縣集集鎮民生東路1號，臺灣。
2. 東海大學生命科學系，40704 臺中市臺中港路3段181號，臺灣。
3. 通信作者。Tel: 886-4-23590121 ext. 32406; Email: slin@thu.edu.tw