



## NOTE

**Notes on *Phalaenopsis* (Orchidaceae) of Hsiaolanyu, Taiwan**Ming-Jer Jung<sup>(1\*)</sup>, Tian-Chuan Hsu<sup>(2)</sup>, Chong-Sheng Leou<sup>(3)</sup> and Ching-Long Yeh<sup>(4)</sup>

1. Institute of Life Science, National Taiwan Normal University, 88, Ting-Chow Rd., Sec 4, Taipei 11677, Taiwan.

2. Herbarium (TAIF), Taiwan Forestry Research Institute, 53, Nanhui Rd., Taipei 10066, Taiwan.

3. Retiree of Experimental Forest of National Taiwan University, 4-5, Hsiaping Rd., Chushan, Nantou 95764, Taiwan.

4. Department of Forestry, National Pingtung University of Science &amp; Technology, 1, Hsuehfu Rd., Neipu, Pingtung 91201, Taiwan.

\* Corresponding Author. Email: l5692401@gmail.com

(Manuscript received 6 January 2010; accepted 11 July 2010)

**ABSTRACT:** Taxonomic position of the *Phalaenopsis* species in Hsiaolanyu, Taiwan is reviewed along with the rediscovery of wild population. *P. riteawanensis* Masam. is lectotypified and reconfirmed to be a synonym of *P. equestris* (Schauer) Rchb.f. Its habitat condition in and conservation status is also presented.

**KEY WORDS:** IUCN category, Lectotypification, Orchidaceae, *Phalaenopsis equestris*, *Phalaenopsis riteawanensis*, Taxonomy, Taiwan.

**INTRODUCTION**

The *Phalaenopsis* species from Hsiaolanyu (a.k.a. Syokotosyo, Ritaiwan and Riteiwan) was first published by Masamune (1934) as a new species, *P. riteiwanensis* Masam. In later studies (Sweet, 1969; Lin, 1977; Su, 2000; Chen and Wood, 2009), *P. riteiwanensis* was treated as a synonym of *P. equestris* (Schauer) Rchb.f. However, the population from Hsiaolanyu was not been collected again since 1934 due to transportation inconvenience and low population size, and its habitat situation and morphological variation are thus nearly uncertain. During a botanical survey of Hsiaolanyu in May 2009, a wild population of *Phalaenopsis* species was rediscovered by our co-worker Sin-Jie Hong and Chuan-Rong Yeh. Later in July 2009, flowers were observed and then confirmed to be *P. riteiwanensis*. The rediscovery of wild population offered us a chance to review the taxonomy position of *Phalaenopsis* species from Hsiaolanyu. After a selection of lectotype and a comparative study on fresh flowering materials, we accept that *P. riteiwanensis* should be a synonym of *P. equestris* (Schauer) Rchb. f., as we present here with a report of habitat condition and conservation status.

**TAXONOMIC TREATMENT**

***Phalaenopsis equestris* (Schauer) Rchb.f.**, Linnaea 22: 864. 1849; Lin, Native Orchids Taiwan 2: 286. f. 44, pl. 140. 1977; Ying, Coloured Ill. Indig. Orchids Taiwan 2: 638. 1990; Tsai et al., Fl. Reipubl. Popularis Sin. 19: 375. 1999; Su, Fl. Taiwan 2nd ed. 5: 1005, 2000; Christenson, *Phalaenopsis*: a Monograph.: 239. 2001; Chen & Wood, Fl. China

25: 382. 2009. Chung, Orchids Taiwan 2: 146. 2009.  
— *Stauroglottis equestris* Schauer, Meyen. Nov. Acad. Cur. 19 (suppl. 1): 432. 1843. — Type: PHILIPPINE. Luzon, Manila, *Lobb s.n.* (holotype: K photo!; isotype: W) 桃紅蝴蝶蘭 Figs. 1-3

*Phalaneopsis riteiwanensis* Masam., Trans. Nat. Hist. Soc. Formosa 24: 213. 1934. — *Stauroglottis riteiwanensis* Masam., Trans. Nat. Hist. Soc. Formosa 24: 213. 1934. pro. *Syn* — Type: TAIWAN. Taitung County, Hsiaolanyu (Syokotosyo; Riteiwan), *K. Segawa s.n.* (lectotype: TAIF no. 115006!, designated here; isolectotypes: TAIF!, KPM photo!).

Distribution and ecology: *Phalaenopsis equestris* is widespread in the Philippines but recorded in Taiwan only from Hsiaolanyu, a small islet located south of Lanyu. Ying (1990) proclaimed that this species also occurred in Lanyu, but there are no specimens available for us to reconfirm this record. Wild population of *P. equestris* in Hsiaolanyu grows in a dwarf forest of ca. 3 m high along the costal cliff of Hsiaolanyu at ca. 100 m alt. The forest is dominated by *Semecarpus gigantifolia* Vidal, *Ficus cumingii* Miq. var. *terminalifolia* (Elm.) Sata, *F. septica* Burm. f. and *Tabernaemontana subglobosa* Merr. Accompanied species include *Planchonella obovata* (R. Br.) Pierre, *Antidesma pentandrum* Merr. var. *barbatum* (Presl.) Merr., *Diospyros ferrea* (Willd.) Bakhuizen, *Pisonia aculeata* L. and *Podocarpus costalis* Presl. All *Phalaenopsis equestris* individuals were found growing on the branches of *Podocarpus costalis*. Flowers were observed from July to August.

Conservation and IUCN category: According to our field investigation and IUCN Red List Categories and Criteria (IUCN, 2001; 2003), we agreed with Lu et al.




**Fig. 1. Lectotype of *Phalaenopsis riteiwanensis* Masam. in the Herbarium of Taiwan Forestry Research Institute (TAIF).**

(2001) and Chung (2009) to consider *Phalaenopsis equestris* Critically Endangered (CR B2ab; D) in Taiwan based on an area of occupancy less than 10 km<sup>2</sup> and matured individuals less than 50. Conservation measures should be taken immediately to ensure the continued existence of *P. equestris* in Taiwan. Although *Phalaenopsis equestris* had been a common species in the Philippines (Lin, 1977), we maintain its regional protection level in Taiwan because the wild population in the Philippines continues deteriorating due to horticultural over-collection (Chung, pers. comm.), and the immigration probability is thus reduced.

**Typification of *Phalaenopsis riteiwanensis*:** After a herbaria reviewing, we found five sheets of *Phalaenopsis* specimens, three in TAIF and two in KPM, fit the type data described by Masamune (1934). Since none of these specimens were marked or defined

as a holotype, lectotypification of *Phalaenopsis riteiwanensis* is necessary for stabilizing its taxonomy position. We select the TAIF no. 115006 specimen as a lectotype because the plant on this sheet is most complete among the possible syntypes (Table 1), and its measurements fit well to the original description of *Phalaenopsis riteiwanensis* (Masamune, 1934).

**Taxonomic Note:** Although *Phalaenopsis equestris* was reported highly variable both vegetatively and in flower colorization (Christenson, 2001), morphology of the population in Hsiaolanyu is quite uniform (Figs. 2 & 3). All individuals we observed bear linear-oblong leaves, strongly reflexed lateral sepals, petals tinged with reddish-purple along midribs, and yellowish-purple lips. The low morphological diversity suggests a single origin of *P. equestris* in Hsiaolanyu.



**Fig. 2. *Phalaenopsis equestris* (Schauer) Reichb. f.** 1: Habit. 2-4: Flower, varied views. 5: Dorsal sepal. 6: Petal. 7: Lateral sepal. 8: Lip. 9: Anther, upper side. 10: Anther, lower side. 11: Column and lip, side view. 12: Lip, longitudinal section. 13: Column, ventral view. 14: Pollinaria. 15 & 16: Pollinia, side views. 17: Stipe with viscidium, upper side. 18: Stipe with viscidium, lower side.

**Table 1. Syntypes of *Phalaenopsis riteawanensis* Masam.**

| Herbarium/Museum |        | Leaves           | Racemose panicle | Flowers | Capsule |
|------------------|--------|------------------|------------------|---------|---------|
| Accession No.    |        |                  |                  |         |         |
| TAIF             | 115006 | 4, 8-18 cm long  | present          | present |         |
|                  | 115007 | absent           | present          | present | absent  |
|                  | 115008 | 2, ca. 6 cm long | present          | present | absent  |
| KPM              | T00076 | absent           | present          | present | absent  |
|                  | T00077 | absent           | present          | present | absent  |



**Fig. 3.** *Phalaenopsis equestris* (Schauer) Rchb. f. from Hsiao Lanyu. A: Plants *in situ*. B-D: Transplanted individual in the greenhouse of National Pingtung University of Science and Technology.

Additional specimens examined: Taiwan. Taitung Co., Hsiao Lanyu, 27 May 1934, S. Sasaki s. n. (TAI), same loc., 24 Aug 2009, M.-J. Jung 4498 (TAIF).

## ACKNOWLEDGEMENTS

We appreciate Dr. Shih-Wen Chung, Dr. Chuan-Rong Yeh and Mr. Sin-Jie Hong for their assistance in field works. We also would like to thank Yi-Shin Chian and Chun-Fen Liao, research assistants of Department of Forestry, National Pingtung University of Science & Technology, Assistant Professor Chih-Wei Chang, Graduate Institute of Marine Biodiversity and Evolutionary Biology, National Dong Hua University and Master Jheng-Jhang Li, research assistants of National Museum of Marine Biology and Aquarium for lots of professional supports. The series of surveys to Hsiao Lanyu was supported in part by grants from Marine National Park Headquarters and National Museum of Marine Biology and Aquarium.

## LITERATURE CITED

- Christenson, E. A. 2001. *Phalaenopsis*, a Monograph. Timber Press, Portland, USA.
- Chen, X.-Q. and J. J. Wood. 2009. *Phalaenopsis*. In: Wu, Z., P. H. Raven and D. Hong (eds.), Flora of China 25: 478-483. Science Press, Beijing, China and Missouri Botanical Garden Press, St. Louis, USA.
- Chung, S.-W. 2008. Orchids of Taiwan 2. Council of Agriculture, Executive Yuan, Taipei, Taiwan.
- IUCN. 2001. IUCN Red List Categories and Criteria, Version 3.1. IUCN Species Survival Commission. IUCN, Gland, Switzerland, and Cambridge, UK.
- IUCN. 2003. Guidelines for Application of IUCN ed List Criteria at Regional Levels, Version 3.0. IUCN Species Survival Commission. IUCN, Gland, Switzerland, and Cambridge, UK.
- Lin, T.-P. 1977. Native Orchids of Taiwan 2. Southern Materials Center, Taipei, Taiwan.
- Lu, S.-Y., S.-J. Moore, T.-H. Hsieh and T.-W. Hsu. 2001. Rare and Endangered Plants in Taiwan 6. Council of Agriculture, Executive Yuan, Taipei, Taiwan.
- Masamune, G. 1934. Beiträge zur Kenntnis der Flora von SüdJapan (II). Trans. Nat. Hist. Soc. Formosa 24: 206-214.
- Su, H.-J. 2000. *Phalaenopsis*. In: Huang, T.-C. et al. (eds.). 2000. Flora of Taiwan. 2nd ed. 5: 1003-1005. Dept. Bot., NTU, Taipei, Taiwan.
- Sweet, H. R. 1969. Revision of the Genus *Phalaenopsis*, pts. 1 and 2. Amer. Orchid Soc. Bull. 37: 867-877, 1089-1104.
- Yeh, C.-L., Y.-S. Chian, C.-F. Liao, C.-R. Yeh, M.-J. Jung and S.-J. Hung. 2010. Inventory of flora in Hsiao Lanyu. Journal of National Park (in press).
- Ying, S.-S. 1990. Coloured Illustrations of Indigenous Orchids of Taiwan 2. Private publication, Taiwan.



## 小蘭嶼蝴蝶蘭屬植物註記

鍾明哲<sup>(1\*)</sup>、許天銓<sup>(2)</sup>、柳重勝<sup>(3)</sup>、葉慶龍<sup>(4)</sup>

1. 國立臺灣師範大學生命科學研究所, 11677 台北市汀州路 4 段 88 號, 臺灣。
2. 林業試驗所植物標本館, 10066 臺北市南海路 53 號, 臺灣。
3. 台大實驗林退休研究員, 95764 南投縣竹山鎮下坪路 4-5 號, 臺灣。
4. 國立屏東科技大學森林系, 91201 屏東縣內埔鄉學府路 1 號, 臺灣。

\* 通信作者。Email: i5692401@gmail.com

(收稿日期：2010 年 1 月 6 日；接受日期：2010 年 7 月 11 日)

**摘要：**隨著蝴蝶蘭屬植物之野生族群於小蘭嶼再度發現，我們重新檢視其分類地位。本文指定了小蘭嶼蝴蝶蘭 (*Phalaenopsis riteiwanensis* Masam.) 之選模式，並再次確認小蘭嶼蝴蝶蘭為桃紅蝴蝶蘭 (*P. equestris* (Schauer) Rchb.f.) 同物異名。本文亦報導桃紅蝴蝶蘭在小蘭嶼的生育地狀態，評估其保育等級。

**關鍵詞：**選模式、蘭科、小蘭嶼蝴蝶蘭、桃紅蝴蝶蘭、分類學、臺灣。