



## NOTE

## ***Frullania evelynae* (Frullaniaceae, Marchantiophyta) – a Very Rare Species from Mizoram, India**

S. K. Singh<sup>(1\*)</sup> and H. A. Barbhuiya<sup>(1)</sup>

1. Botanical Survey of India, Eastern Regional Centre, Shillong – 793003, India.

\* Corresponding author. Email: sksbsinc@rediffmail.com

(Manuscript received 25 February 2011; accepted 15 March 2011)

**ABSTRACT:** *Frullania evelynae* S. Hatt. & Thaith.- a very rare species of family Frullaniaceae of Hepaticae is recorded from Mizoram, India for the first time. A detailed taxonomic description along with line drawing plate is provided for its easy identification. A note on diversity and distribution of Indian *Frullania*'s which is in chaos in recent past is also provided.

**KEY WORDS:** *Frullania evelynae*, Hepaticopsida, Mizoram, India.

### **INTRODUCTION**

The genus *Frullania* Raddi (Frullaniaceae) is represented by ca. 1000 species in the world (Yuzawa, 1991). In India the genus is represented by approximately 66 taxa incl. 3 subspecies, 8 varieties and 2 forma (Mitten, 1861; Stephani, 1910; Verdoorn, 1930; Kashyap, 1932; Chopra, 1943; Hattori, 1966, 1971, 1972, 1973, 1974, 1975, 1978, 1980, 1980a, 1981, 1982, 1986; Hattori and Lin 1985; Hattori and Thaithong, 1978, 1978a, 1978b; Srivastava, 1979; Lal, 1979; Lal and Parihar, 1979; Udar and Nath, 1981; Udar and Kumar, 1983; Nath, 1984; Nath and Udar, 1984; Joshi et al., 1989; Nath and Asthana, 1992, 1998; Parihar et al., 1994; Bapna and Kachroo, 2000; Srivastava and Alam, 2002, 2004; Singh and Nath, 2004, 2007; Singh and Singh 2003, 2005, 2007; Nath and Singh 2006; Dey and Singh, 2008; Dey et al., 2009; Daniels, 2010), distributed mainly in Eastern Himalaya, Western Himalaya and Western Ghats (Table 1).

*Frullania evelynae* belongs to the subgenus *Trachycolea* Spruce was described originally by Hattori and Thaithong in 1978 on the basis of collections made by Sharp and Iwatsuki from Khasi & Jaintia Hills and Darjeeling area in India. This species never reported either from India or elsewhere except by Sinske Hattori and P.J. Lin from Yunnan, China (Hattori and Lin, 1985). However, the area was thoroughly surveyed by many bryologists in recent past but could not be collected from any locality including its type locality. Recently, a comprehensive work has been done and Floristic account was published by Singh and Nath on *Hepaticae of Khasi and Jaintia Hills* (Singh and Nath, 2007) but they could describe this species on the basis of its type collection only.

Recently, during a field exploration to document the Bryoflora (Hepaticae and Anthocerotae) of Mizoram, in 2008, the senior author came across with a small epiphytic population of this species. Critical study of the plants followed by scrutiny of literature reveals it's identity as *Frullania evelynae*. It is interesting to note that this species is new record for State of Mizoram and also constitutes the second collection from India after its Type.

### **TAXONOMIC TREATMENTS**

***Frullania evelynae* S. Hatt. & Thaith. in J. Hattori Bot. Lab. 44: 179. 1978; S. Hatt. & P. J. Lin in J. Hattori Bot. Lab. 59: 134. 1985. (Fig. 1: 1-21.)**

Plants light green, 35-45 mm long, 1.4-2.0 mm wide, irregularly 1-2 pinnately branched. Stem in cross-section spherical in outline, 0.4-0.5 mm, 10-12 cells across, not differentiated into cortex and medulla; cells triangular to subquadangular or globose, 17.5-35.0 × 12.5-25.0 µm. Leaves complicate-bilobed, leaf-lobe ovate to nearly orbicular, imbricate, wide spreading, dorsally extending beyond the farther edge of stem, slightly concave, 1.2-1.5 mm long, 1.0-1.1 mm wide, more or less incurved at apex, marginal cells towards apex quadrangular, 12.5-20 × 12.5-17.5 µm, walls slightly flexuose, median cells subquadangular to polygonal, 20-30 × 15-25 µm, with incrassate and confluent trigones, basal cells pentagonal, 20-50 × 17.5-30 µm, with nodulose-bulging trigones, intermediate thickening gives impression of sinuate thickenings; leaf lobule cucullate, contiguous to the stem, truncate, with incurved beak extending slightly downward beyond the ventral margin, 0.3-1.0 mm long,

**Table 1. Distribution of genus *Frullania* in India**

Species name	EH	WH	WG	CI	A&N	References
1. <i>F. acutiloba</i> Mitt.	+		+			Mitten, 1861; Hattori & Lin, 1985; Nath & Asthana, 1998; Srivastava & Alam, 2002; Singh & Nath, 2007; Daniels, 2010 [Hattori, 1973 as <i>F. asperula</i> Mitt.; Hattori, 1975a as <i>F. hampeana</i> var. <i>acutiloba</i> (Mitt.) S.Hatt.]
2. <i>F. alstonii</i> Verd. var. <i>pfleidereri</i> S. Hatt.	+					Hattori & Thaithong, 1978; Singh & Nath, 2007
3. <i>F. apiculata</i> (Reinw. et al.) Dumort.	+		+		+	Mitten, 1861; Hattori & Thaithong, 1978; Nath & Asthana, 1998; Singh & Nath, 2007; [Hattori & Thaithong, 1978a as <i>F. apiculata</i> (Reinw. et al.) Dumort. var. <i>obtusata</i> S.Hatt. & Thaith.]
4. <i>F. arecae</i> (Spreng.) Gottsche	+	+	+			Yuzawa, 1991; Daniels, 2010 [Stepahni, 1910 as <i>F. indica</i> Steph; Mitten, 1861; Hattori, 1966, 1975; Nath & Asthana, 1998; Srivastava & Alam 2002; Singh & Nath, 2007 as <i>F. wallichiana</i> Mitt.]
5. <i>F. campanulata</i> Sande Lac.				+		Nath & Asthana, 1998; Srivastava & Alam, 2002
6. <i>F. debilis</i> Steph. ex S. Hatt.				+		Hattori, 1974; Daniels, 2010
7. <i>F. davurica</i> Hampe		+				Hattori & Lin, 1985 [Singh & Nath, 2004, 2007 as <i>F. rotudistipula</i> Steph.]
8. <i>F. densiloba</i> Steph.						Srivastava & Alam, 2004
9. <i>F. dialatata</i> (L.) subsp. <i>asiatica</i> S. Hatt.	+	+				Hattori, 1982; Hattori & Thaithong, 1978b
10. <i>F. dialatata</i> (L.) subsp. <i>dilatata</i>			+			Singh & Singh, 2007
11. <i>F. duthiana</i> Steph. var. <i>duthiana</i>	+	+				Stephani, 1910; Hattori, 1966; Hattori & Thaithong, 1978; Nath & Udar, 1984
12. <i>F. duthiana</i> Steph. var. <i>appendiculata</i> S. Hatt.	+					Hattori, 1973, 1978
13. <i>F. ericoides</i> (Nees) Mont.	+	+	+	+	+	Hattori, 1971; Hattori & Thaithong, 1978, 1978a; Bapna & Kachroo, 2000 [Hattori, 1966; Nath & Asthana, 1998; Srivastava & Alam, 2002; Singh & Nath, 2007 as <i>F. squarrosa</i> ; Singh & Singh, 2003 as <i>F. squarrosa</i> f. <i>ericoides</i> (Verd.) Nees; Mitten, 1861 as <i>F. squarrosa</i> and also <i>F. ericoides</i> ]
14. <i>F. evelynae</i> S. Hatt. & Thaith.	+					Hattori & Thaithong, 1978; Nath & Udar, 1984
15. <i>F. evoluta</i> Mitt.	+		+			Hattori, 1974, 1975; Daniels, 2010
16. <i>F. gaudichaudii</i> (Nees & Mont.) Nees & Mont.			+		+	Nath & Asthana, 1998; Joshi et al., 1989; Hattori, 1966, 1972; Hattori & Lin, 1985 [Mitten 1961 as <i>F. recurvata</i> Lehm. & Lindenb. and also as <i>F. ceylanica</i> Nees]
17. <i>F. gemmulosa</i> S. Hatt. & Thaith.	+					Hattori & Thaithong, 1978
18. <i>F. gracilis</i> (Reinw. et al.) Dumort.					+	Hattori, 1974
19. <i>F. gracilis</i> (Reinw. et al.) Dumort. subsp. <i>zennoskei</i> . & Thaith.	+					Hattori & Thaithong, 1978
20. <i>F. geraldiana</i> Massal.	+					Singh & Nath, 2007
21. <i>F. grandistipula</i> Lindb.	+	+	+			Hattori, 1986; Nath, 1984; Singh & Nath, 2007
22. <i>F. hattoriantha</i> Udar & Nath	+					Udar & Nath, 1981; Dey et al., 2009
23. <i>F. himalayensis</i> Steph.			+			Hattori & Thaithong, 1978; Hattori, 1978
24. <i>F. inflexa</i> Mitt.	+		+			Mitten, 1861; Stephani, 1910; Hattori, 1975; Hattori & Thaithong, 1978; Nath & Asthana, 1998 [Hattori, 1966, 1971 as <i>F. delavayi</i> Steph.]
25. <i>F. inftata</i> Gottsche	+	+				Hattori & Lin, 1985; Hattori & Thaithong, 1978 [Parihar et al, 1994 as <i>F. mayebarae</i> S. Hatt.]
26. <i>F. inftata</i> Gottsche var. <i>dioica</i> S. Hatt. & Thaith.			+			Hattori & Thaithong, 1978
27. <i>F. intermedia</i> (Reinw. et al.) Dumort.					+	Joshi et al., 1989; Nath & Asthana, 1998; Hattori, 1980a
28. <i>F. larjana</i> Sushil K. Singh & D. K. Singh			+			Singh & Singh, 2005
29. <i>F. meyeniana</i> Lindenb.	+					Hattori & Thaithong, 1978
30. <i>F. monocera</i> (Tayl.) Tayl.	+					Bapna & Kachroo, 2000 [Robinson, 1964 as <i>F. hampeana</i> Nees]
31. <i>F. montana</i> Steph.	+					Hattori & Thaithong, 1978

**Table 1. Continued.**

Species name	EH	WH	WG	CI	A&N	References
32. <i>F. muscicola</i> Steph.	+	+	+	+		Robinson, 1964; Srivastava, 1979; Kashyap, 1932; Hattori, 1966; Hattori & Thaithong, 1978; Nath & Asthana, 1998; Bapna & Kachroo, 2000; Narayan et al., 2001; Srivastava & Alam, 2002; Singh & Nath 2007
33. <i>F. muscicola</i> Steph. var. <i>inaeana</i> (Steph.) Kamim.	+			+		Lal & Parihar, 1979; Dey et al., 2009
34. <i>F. nepalensis</i> (Spreng.) Lehm. & Lindenb.	+	+				Mitten, 1861; Hattori, 1966; Nath, 1977; Hattori & Thaithong, 1978, 1978a [Hattori, 1966, 1971 also as <i>F. grevilleana</i> Taylor]
35. <i>F. nepalensis</i> (Spreng.) Lehm. & Lindenb. f. <i>amphigastriis</i> Mitt.	+	+				Mitten, 1861; Chopra, 1943; Hattori, 1966.
36. <i>F. neurota</i> Taylor	+	+	+			Mitten, 1861; Hattori, 1966, 1971; Yuzawa, 1991; Nath & Asthana, 1992; Singh & Nath, 2007; [Mitten 1861 also as <i>F. breviuscula</i> Mitt.]
37. <i>F. neurota</i> Taylor var. <i>pfeiffereri</i> S. Hatt.	+					Hattori, 1972
38. <i>F. nodulosa</i> (Reinw. et al.) Nees			+	+		Hattori, 1980a; Daniels, 2010
39. <i>F. pallide-virens</i> Steph.	+					Hattori & Thaithong, 1978; Dey et al., 2009
40. <i>F. pariharii</i> S. Hatt. & Thaith.	+		+			Hattori & Thaithong, 1978, 1978a
41. <i>F. pariharii</i> S. Hatt. & Thaith. f. <i>intermedia</i> Thaith.	+					Hattori & Thaithong, 1978
42. <i>F. physantha</i> Mitt.	+					Mitten, 1861; Stephani, 1910; Hattori, 1966, 1971; Hattori & Thaithong, 1978; Nath & Udar, 1984
43. <i>F. pran-nathii</i> M. Dey & D. K. Singh	+					Dey & Singh, 2008.
44. <i>F. polyptera</i> Taylor		+	+			Hattori, 1980; Hattori & Lin, 1985 [Kashyap, 1932 as <i>F. pyriflora</i> Steph. also as <i>F. gollani</i> Steph.; Mitten, 1861 as <i>F. subinflata</i> Mitt.]
45. <i>F. pseudoschensiana</i> S. Hatt.	+					Hattori, 1980; Dey et al., 2009
46. <i>F. pseudoschensiana</i> S. Hatt. var. <i>darjeelingensis</i> S. Hatt.	+					Hattori, 1981; Singh & Nath, 2007
47. <i>F. ramuligera</i> (Nees) Mont.			+			Daniels, 2010
48. <i>F. retusa</i> Mitt. var. <i>retusa</i>	+	+				Chopra, 1943; Hattori, 1966, 1971; Hattori & Thaithong, 1978a; Nath & Udar, 1984
49. <i>F. retusa</i> Mitt. var. <i>hirsuta</i> S. Hatt. & Thaith.	+	+				Hattori & Thaithong, 1978
50. <i>F. retusa</i> Mitt. var. <i>gymnantha</i> S. Hatt. & Thaith.		+				Hattori & Thaithong, 1978a
51. <i>F. rhytidanta</i> S. Hatt.	+					Hattori, 1980
52. <i>F. riojaneirensis</i> (Raddi) Spruce			+			Srivastava & Alam, 2004
53. <i>F. serrata</i> Gottsche f. <i>serrata</i>	+		+			Verdoorn, 1930; Hattori, 1973; Nath & Asthana, 1998; Singh & Nath, 2007
54. <i>F. schensiana</i> Massal.	+					Hattori, 1966, 1971
55. <i>F. sphaerantha</i> S. Hatt.						Hattori, 1980
56. <i>F. sharpantha</i> Udar & Kumar	+					Udar & Kumar, 1983
57. <i>F. shanensis</i> Svhila	+					Hattori & Thaithong, 1978b
58. <i>F. sinensis</i> Steph.	+	+				Hattori, 1980; Mitten, 1861; Chopra, 1943; Bapna & Kachroo, 2000
59. <i>F. subclavata</i> Steph.	+	+				Hattori, 1975, 1980; Hattori & Thaithong, 1978, 1978a
60. <i>F. subpedicellata</i> S. Hatt.	+					Hattori, 1980
61. <i>F. tamarisci</i> (L.) Dumort.	+		+			Nath & Asthana, 1998; Srivastava & Alam, 2002; Singh & Nath, 2007; Dey et al., 2009 [Mitten, 1861 as <i>F. moniliata</i> Nees]
62. <i>F. tamarisci</i> (L.) Dumort. subsp. <i>obscura</i> (Verd.) S. Hatt.	+					Hattori & Thaithong, 1978
63. <i>F. tagawana</i> (S. Hatt. & Thaith.) S. Hatt.	+					Hattori & Lin, 1985 [Hattori & Thaithong, 1978 as <i>F. evoluta</i> Mitt. var. <i>tagawana</i> S. Hatt. & Thaith.]
64. <i>F. tuyamae</i> S. Hatt. & Thaith.	+					Lal, 1979
65. <i>F. udarii</i> Nath & A. P. Singh	+					Nath & Singh, 2006
66. <i>F. yunnanensis</i> Steph.	+	+				Hattori, 1966, 1971; Hattori & Thaithong, 1978; Nath & Udar, 1984

EH: Eastern Himalaya; WH: Western Himalaya; WG: Western Ghats, CI: Central India; A &amp; N: Andaman and Nicobar Islands.




Fig. 1. *Frullania evelynae* S. Hatt. & Thaith. 1: A portion of the plant in dorsal view. 2: The same in ventral view. 3: C.S. of stem. 4-6: Leaves with lobule. 7: Marginal leaf cells towards apex. 8: Median leaf cells. 9: Basal leaf cells. 10-11: Styls. 12-15: Underleaves. 16: Female inflorescence showing arrangement of bract bracteole and perianth. 17: Female bract. 18: Female bract with connate bracteole. 19-20: Female bracteoles. 21: C.S. perianth.

0.15-0.5 mm wide; styls minute, usually 6-11celled, cells arranged in a row, sometimes at base up to 2 cells wide, 87.5-142.5  $\mu$ m long, 7.5-37.5  $\mu$ m. Underleaves approximate to imbricate towards apex, suborbicular to

reniform, entire, 0.6-0.9 mm long, 0.8-1.2 mm wide, 2-2.5 times the width of stem, nearly flat, margin slightly undulated and incurved. Male inflorescence not seen. Female inflorescence terminal on main stem or



branches. Perianth semi-exerted, pyriform, 1.7-2.0 mm long, 1.0-1.2 mm wide, 5-keeled, keels high and narrow, undulated, surface smooth, apex obtuse with short apical beak; female bracts in 3 pairs, bract lobe of inner whorl ovate-orbicular with subacute apex, 1.4-2.0 mm long, 1.0-1.2 mm wide; inner bract lobule 1/2 -connate, ovate, with acute apex, 0.7-1.8 mm long, 0.5-0.9 mm wide, outermost bract lobule sometimes lanceolate-canaliculate ca. 0.7 mm long, 0.5 mm wide; bracteoles broadly ovate or rectangular, margin strongly incurved 0.8-1.7 mm long, 0.4-1.2 mm wide, 1/6-1/5<sup>th</sup> bifid, lobes triangular with subacute apex.

**Specimen examined:** Mizoram, Lungtai, Towads S. Vataphaphai, growing on tree trunk in association with *Cheilolejeunea* sp., 29.06.2008. S. K. Singh 119888 (ASSAM).

**Distribution:** INDIA: Meghalaya (Khasi & Jaintia Hills); West Bengal (Darjeeling); Mizoram (-present study). CHINA.

*Frullania evelynae* is characterized by entire suborbicular to reniform underleaves and semi-exerted, pyriform with 5 high, narrow, undulated keeled perianth. This species is closely allied with *F. physntha* Mitt. However, the latter is clearly distinct from the former by having large exerted perianth which are almost spherical, somewhat prominulate at apex with 5 short, more or less angular-crispate keels and the innermost female bract with an acute lobe and acuminate-acute lobule bearing 1 large tooth or lacina on the ventral margin and the innermost female bracteole with 2 apical teeth and often with minute basal teeth (Thaithong and Hattori, 1977).

#### Note on diversity and distribution of *Frullania* in India

So far about 66 taxa of *Frullania* recorded from different Bryogeographical regions of India. Among them, with 50 taxa Eastern Himalaya is richest followed by Western Himalaya with 21 taxa and Western Ghats with 19 taxa. Andaman & Nicobar Islands is represented with 6 taxa and Central India with 3 taxa. Whereas the other bryological territories has no representation. Out of 66 taxa, 14 taxa viz., *F. debilis* Steph. ex S. Hatt., *F. gracilis* (Reinw. et al.) Dumort. subsp. *zennoskei* S. Hatt. & Thaith., *F. hattoriantha* Udar & Nath, *F. inflata* Gottsche var. *dioica* S. Hatt. & Thaith., *F. larjiana* Sushil K. Singh & D. K. Singh, *F. pariharii* S. Hatt. & Thaith. f. *intermedia* Thaith. *F. pran-nathii* M. Dey & D. K. Singh, *F. pseudoscheniana* S. Hatt., *F. retusa* Mitt. var. *hirsuta* S. Hatt. & Thaith. *F. retusa* Mitt. var. *gymnantha* S. Hatt. & Thaith., *F. rhytidanta* S. Hatt., *F. sphaerantha* S. Hatt., *F. subpedicellata* S. Hatt. *F. udarii* Nath & A. P. Singh are endemic to Indian regions (Table 1).

#### Doubtful species

*F. glomerata* (Lehm. & Lindenb.) Mont. – listed from the Nilgiri Mts. by Montagne (1842) and Mitten (1861), but the species is confined to tropical C. and S. America (Gradstein & Costa 2003). It is not known to which taxon the Indian specimens should be assigned.

*F. riparia* Hampe ex Lehm. (syn. *F. aelotis*) – confined to Europe eastwards to Iran, and North America, Mexico (Schuster, 1992).

#### ACKNOWLEDGEMENTS

Authors are grateful to Director, Botanical Survey of India, Kolkata and to Head of the office, BSI, Eastern Regional Centre, Shillong for facilities and encouragements and to the officials of Forest Department of Mizoram for rendering the logistic support during field exploration.

#### LITERATURE CITED

- Bapna, K. R and P. Kachroo.** 2000. Hepaticology in India. Himanshu Publication Udaipur & New Delhi, India.
- Chopra, R. S.** 1943. A census of Indian Hepaticas. J. Indian Bot. Soc. **22**: 237-259.
- Daniels, A. E. D.** 2010. Checklist of the bryophytes of Tamil Nadu, India. Arch. Bryol. **65**: 1-117.
- Dey, M. and D. K. Singh.** 2008 *Frullania pran-nathii* – a new epiphyllous liverwort from Darjeeling, India. J. Jap. Bot. **83**: 280-283.
- Dey, M., D. Singh and D. K. Singh.** 2009. Some new and noteworthy records of Hepaticae from eastern Himalaya, India. Indian J. Forest. **16**: 669-684.
- Gradstein, S. R. and D. P. Da Costa.** 2003. The Hepaticae and Anthocerotae of Brazil. Mem. New York Bot. Gard. **87**: 1-318.
- Hattori, S.** 1966. Hepaticae and Anthocerotae. In Hara, H. (ed.) The flora of Eastern Himalaya. University of Tokyo, Tokyo pp. 501-586.
- Hattori, S.** 1971. Hepaticae. In Hara, H. (ed.) The flora of Eastern Himalaya. Second report. University of Tokyo Press, Tokyo. pp. 222-240.
- Hattori, S.** 1972. Notes on Asiatic species of the genus *Frullania*, Hepaticae. I. J. Hattori Bot. Lab. **36**: 109-140.
- Hattori, S.** 1973. Notes on Asiatic species of the genus *Frullania*, Hepaticae. II, III, IV. J. Hattori Bot. Lab. **37**: 55-152.
- Hattori, S.** 1974. Notes on Asiatic species of the genus *Frullania*, Hepaticae. V, VI. J. Hattori Bot. Lab. **38**: 185-274.
- Hattori, S.** 1975. A revision of Indian species of *Frullania* (Hepaticae) published by W. Mitten in 1861. Bull. Natl. Sci. Mus. Ser B. **1**: 73-81.
- Hattori, S.** 1975a. Notes on Asiatic species of the genus *Frullania*, Hepaticae. VII. J. Hattori Bot. Lab. **39**: 277-313.
- Hattori, S.** 1978. Notes on the Asiatic species of the genus *Frullania*, Hepaticae. XI. J. Hattori Bot. Lab. **44**: 525-554.
- Hattori, S.** 1980. Notes of the Asiatic species of the genus *Frullania*, Hepaticae. XII. J. Hattori Bot. Lab. **47**: 85-125.



- Hattori, S.** 1980a. A revision of the subgenus *Homotropantha* of the genus *Frullania*, Hepaticae. J. Hattori Bot. Lab. **47**: 165-236.
- Hattori, S.** 1981. Notes of the Asiatic species of the genus *Frullania*, Hepaticae. XIII. J. Hattori Bot. Lab. **47**: 147-168.
- Hattori, S.** 1982. Asian taxa of the *Frullania dilatata* complex. J. Jap. Bot. **57**: 257-260.
- Hattori, S.** 1986. Notes of the Asiatic species of the genus *Frullania*, Hepaticae. XIV. Bull. Nat. Sci. Mus. Ser. B. **12**: 177-138.
- Hattori, S. and P. J. Lin.** 1985. A preliminary study of Chinese *Frullania* flora. J. Hattori Bot. Lab. **59**: 123-169.
- Hattori, S. and O. Thaithong.** 1978. Indian *Frullania* collection made by Prof. & Mrs. A. J. Sharp and Dr. Z. Iwatsuki. J. Hattori Bot. Lab. **44**: 177-193.
- Hattori, S. and O. Thaithong.** 1978a. Dr. Parihar's collection of Indian *Frullania* (Hepaticae). J. Jap. Bot. **53**: 129-133.
- Hattori, S. and O. Thaithong.** 1978b. Mrs. Ruth D. Svihla's *Frullania* collections in India, Burma and Singapore. Bull. Natl. Sci. Mus. Ser. B **4**: 65-69.
- Joshi, D. Y., D. D. Wani and S. J. Chavan.** 1989. Studies on the hepatic flora of the Andaman islands (India)-I. Ad. Pl. Sci. **2**: 169-177.
- Kashyap, S. R.** 1932. Liverworts of the Western Himalayas and the Panjab Plain. II. University of the Punjab, Lahore.
- Lal, J.** 1979. *Frullania tuyamae* Hatt. & Thaith. (Hepaticae) from Manipur (eastern India)-new to India. Bull. Bot. Surv. India. **21**: 81-83.
- Lal, J. and N. S. Parihar.** 1979. Contributions to the Bryoflora of Central Indian Zone I-liverworts. J. Indian Bot. Soc. **58**: 110-114.
- Mitten, W.** 1861. Hepaticae Indiae Orientalis, an enumeration of the Hepaticae of the East Indies. J. Proc. Linn. Soc. Bot. **5**: 87-108.
- Montagne, J. P. F. C.** 1842. Cryptogamae Nilgherienses Plantarum cellularium in montibus peninsulae Indicae Neel-Gherries dictis à cl. Perrottet collectarum Enumeratio, Hepaticae. Ann. Sci. Nat., Bot. 2, **18**: 12-17.
- Narayan, V., P. V. Karunakaran and D. K. Singh.** 2001. Contribution to the Bryoflora of Great Himalayan National Park, Kullu, Himachal Pradesh-1. Indian J. Forest. **24**: 265-278.
- Nath, V.** 1984. Abnormal elaters in *Frullania grandistipula* Ldbg.- a new record. Nat. Acad. Sci. Lett. **1**: 297-298.
- Nath, V. and A. K. Asthana.** 1992. Addition to the Bryoflora of Western Himalaya: *Frullania neurota* Taylor. Geophytology **21**: 139-141.
- Nath, V. and A. K. Asthana.** 1998. Diversity and distribution of genus *Frullania* Raddi in South India. J. Hattori Bot. Lab. **85**: 63-82.
- Nath, V. and A. P. Singh.** 2006. *Frullania udarei* Nath et Singh - A new species from Meghalaya, India. Curr. Sci. **91**: 744-746.
- Nath, V. and R. Udar.** 1984. Genus *Frullania* Raddi in India I. Species with entire amphigastria. Proc. Indian Natl. Sci. Acad. B **50**: 223-234.
- Parihar, N. S., B. Lal and N. Katiyar.** 1994. Hepaticae and Anthocerotae of India. A new annotated checklist. Allahabad Central Book Depot. Allahabad, India.
- Robinson, H.** 1964. A small collection of Bryophytes from Upper Assam. J. Hattori Bot. Lab **27**: 124-130.
- Schuster, R. M.** 1992. The Hepaticae and Anthocerotae of North America, Vol. V. Field Museum of Natural History, Chicago
- Singh, A. P. and V. Nath.** 2004. *Frullania rotundistipula* Steph. - an addition to the Indian bryoflora. Geophytology **32**: 73-77.
- Singh, A. P. and V. Nath.** 2007. Hepaticae of Khasi and Jaintia Hills: Eastern Himalayas. Bishen Singh Mahandrapal Singh, Dehradun, India. 382pp.
- Singh, A. P., D. Kumar and V. Nath.** 2008. Studies on the Genera *Frullania* Raddi and *Jubula* Dum. from Meghalaya (India): Eastern Himalayas. Taiwania **53**: 51-84.
- Singh, S. K. and D. K. Singh.** 2003. Contribution to the Bryoflora of Great Himalayan National Park, Kullu, Himachal Pradesh, India - II. Hepaticae. Phytotaxonomy **3**: 35-52.
- Singh, S. K. and D. K. Singh.** 2005. *Frullania larjiana* (Jungermanniopsida: Frullaniaceae), a new species from India. J. Bryol. **27**: 105-108.
- Singh, S. K. and D. K. Singh.** 2007. Some new and noteworthy records of Hepaticae and Anthocerotae from Western Himalaya, India. Cryptog. Bryol. **28**: 253-265.
- Srivastava, S. C.** 1979. Hepaticae of Kashmir Valley. Nova Hedwigia **63**: 333-338.
- Srivastava, S. C. and A. Alam.** 2002. A collection of *Frullanin* from Nilgiri with *F. densiloba* St. as a new record for India. J. Bombay Nat. Hist. Soc. **99**: 232-237.
- Srivastava, S. C. and A. Alam.** 2004. *Frullania riojaneirensis* (Raddi) Spruce - an addition to Indian bryoflora. Geophytology **33**: 47-51.
- Stephani, F.** 1910. Species Hepaticarum vol. 4. Geneva.
- Thaithong, O. and S. Hattori.** 1977. A new species of *Frullania* From N. Vietnam and the Indian *F. physantha* Mitt. Bull. Natl. Sci. Mus. Bot. **3**: 149-154.
- Udar, R. and A. Kumar.** 1983. A new *Frullania* from India. Misc. Bryol. Lechenol. **9**: 192-194.
- Udar, R. and V. Nath.** 1981. *Frullania hattoriantha* sp. nov. from India. Misc. Bryol. Lechenol. **9**: 44-47.
- Verdoorn, F.** 1930. Die Frullaniaceae der Indomalasis Chen Insulin (Die Frullaniacis VII). Ann. Bryol. (supp.) **1**: 181-187.
- Yuzawa, Y.** 1991. A monograph of subgen. *Chonanthelia* of genus *Frullania* (Hepaticae) of the world. J. Hattori Bot. Lab. **70**: 181-291.



## 伊氏耳葉蘚 (*Frullania evelynae*, 耳葉蘚科, 地錢門) – 印度米佐蘭姆的一個稀有種

S. K. Singh<sup>(1\*)</sup> and H. A. Barbhuiya<sup>(1)</sup>

1. Botanical Survey of India, Eastern Regional Centre, Shillong – 793003, India.

\* 通信作者。Email: sksbsinc@rediffmail.com

(收稿日期：2011年2月25日；接受日期：2011年3月15日)

**摘要：**在印度米佐蘭姆 (Mizoram) 首次記錄到伊氏耳葉蘚 (*Frullania evelynae*) 這個非常罕見的耳葉蘚科 (Frullaniaceae) 物種。本文提供了詳細分類說明及手繪圖以利辨識；並提供印度地區耳葉蘚屬 (*Frullania*) 物種的清單及分佈。

**關鍵詞：**伊氏耳葉蘚、苔綱、印度、米佐蘭姆。