



NOTE

Thuidium recognitum (Hedw.) Lindb. var. *delicatulum* (Hedw.) Warnst – a Moss New to Indian Bryoflora

Virendra Nath^(1*), Pooja Bansal⁽¹⁾ and A. K. Asthana⁽¹⁾

1. Bryology Laboratory, National Botanical Research Institute (Council of Scientific and Industrial Research, New Delhi, India) Lucknow – 226 001, India.

* Corresponding author. Tel.: 0522- 2297837; Fax: 0522- 2205836, 2205839; Email: drvirendranath2001@rediffmail.com

(Manuscript received 7 April 2011; accepted 23 May 2011)

ABSTRACT: *Thuidium recognitum* belonging to Musci under family Thuidiaceae is reported for the first time from India. The specimens collected on the way to Dikhu river, Mokokchung, North Eastern hills, Nagaland highly resemble with the variety of *T. recognitum* var. *delicatulum* in acute stem leaf with top conical leaf papillae, dentate costa ending below tip. It is an addition to bryoflora of India, but has its restricted distribution in eastern Himalaya.

KEY WORDS: Eastern Himalaya, India, moss, new record, Thuidiaceae, *Thuidium*.

INTRODUCTION

Genus *Thuidium* Bruch, Schimper & Gumbel, a pleurocarpous moss of moist temperate region belonging to family Thuidiaceae was established by Bruch *et al.* (1836-1866). It is a cosmopolitan genus with about 158 valid species all over the world (Gangulee, 1978). As far as number of species pertaining to this genus in India is concerned, different workers have different opinions. Robinson (1968) reported 16 species of this genus from Himalaya, whereas Gangulee (1978) reported 18 species from various part of the country. Later on Vohra (1983) reported 20 species from India. Watanabe (1991) reported 14 species of this genus from Asia, of which 12 are from India. Recently, Lal (2005) provided a checklist of Indian mosses and reported 24 species of this genus from India.

Revisionary work on *Thuidium* has been carried out in Japan and adjacent regions (Watanabe, 1972), South of the Sahara Desert (Touw, 1976, 2001), Western Melanesia (= Eastern Malesia) (Norris and Koponen, 1985), Australasia (Touw and Haak, 1989) and North America (Crum, 1999). In India, Gangulee (1978) synonymised *T. delicatulum* under *T. recognitum* (Hedw.) Lindb. var. *delicatulum* (Hedw.) Warnst, also, Kuo and Chiang (1987) and Noguchi *et al.* (1991) retained the status of *T. recognitum* var. *delicatulum* as such and synonymised *T. delicatulum* under it. However, Kürschner and Erdağ (2005) treated *Thuidium delicatulum* (Hedw.) Mitt. and *T. recognitum* as separate species.

In this paper, the authors followed the concept of

Gangulee (1978), Kürschner and Erdağ (2005), Kuo and Chiang (1987) and Noguchi *et al.* (1991), and described *delicatulum* at varietal level of *T. recognitum*. It is a new record for the bryoflora of India, though restricted in eastern Himalaya.

MATERIALS AND METHODS

The plants grow on moist soil on the way to Dikhu river, Mokokchung, Nagaland, and they were collected in August, 2008 associated with *Hypnum subimponens* Lesq. ssp. *ulophyllum* (C. Muell.) Ando and *Leucobryum nilghiriense* C. Muell. The specimens have been deposited in the Bryophyte Herbarium, National Botanical Research Institute, Lucknow (LWG).

TAXONOMIC TREATMENT

Thuidium recognitum (Hedw.) Lindb. var. *delicatulum* (Hedw.) Warnst, Bot. Centralbl., 5: 185 (1881).

Fig. 1

Hypnum delicatulum Hedw., *Spec. Musc.*: 260 (1801); *Thuidium delicatulum* (Hedw.) Mitt., *J. Linn. Soc. Bot.*, 12: 578 (1869) nom. illeg.; *Thuidium viride* Mitt., *Trans. Linn. Soc. London Bot. ser. 2, 3*: 188 (1891); *Thuidium uliginosum* Card., *Bull. Soc. Bot. Genève, sér. 2, 3*: 283 (1911); *Thuidium planopteris* Sak., *Bot. Mag. Tokyo*, 57: 349 (1943); *Thuidium tetracladioides* Sak., *Bot. Mag. Tokyo*, 60: 89 (1947); *Thuidium yezoanum* Sak., *Bot. Mag. Tokyo*, 60: 89 (1947); *Thuidium iyoense* Sak. in Oti, *Bot. Mag. Tokyo*, 63: 81 (1950); *Thuidium xanthobasis* Sak., *Bot. Mag. Tokyo*, 64: 199 (1951); *Thuidium tetracladioides* var. *xanthobasis* (Sak.) T. Inoue, *Misc. Bryol. Lichenol.* 12: 4 (1957); *Thuidium recognitum* auct. non. (Hedw.) Lindb., Noguchi, *J. Hattori Bot. Lab.* 20: 285 (1958).




Fig. 1. *Thuidium recognitum* (Hedw.) Lindb. var. *delicatulum* (Hedw.) Warnst. A: Plant (diagrammatic). B: Habit. C: Portions of plant. D: Cross section of stem. E: Paraphyllia. F-J: Stem leaves. K-R: Branch leaves. S: Apical cells of stem leaf. T: Median cells of stem leaf. U: Median cells of branch leaf. V: Basal cells of branch leaf.



Plants large, ±robust, forming light – green to yellowish to brownish mats. Stems prostrate, creeping, up to 12 cm long, bipinnately branched. Paraphyllia abundant at or near cell ends, papillose, the terminal cell usually acuminate, rarely obtuse or bifurcate, with 2-3 large, divergent papillae. Stem leaves erect when dry, widely ovate – lanceolate, $1.05-1.32 \times 0.67-0.72$ mm, abruptly acuminate. Margin sharply toothed. Costa stout, ending in acumen below tip, dentate. Cells pellucid, 12.5×8.3 μm , unipapillose. Few rows of cells at base larger, rectangular, 16.7×8.34 μm . Branch leaves dense, erectopate, concave, ovate – lanceolate, 0.5×0.3 mm, with acute tip. Leaf margin crenulate, costa ending below tip, leaf cells with single and sharp papillae. Sporophytes not seen.

Specimens examined: India: Eastern Himalaya, Nagaland, Mokokchung, on way to Dikhu river, 06.08.2008, Coll. by: V. Nath, 248857B, 248858B (LWG).

DISCUSSION

T. recognitum (Hedw.) Lindb. var. *delicatulum* (Hedw.) Warnst is allied to *T. orientale* Mitt. ex Dix. in most morphological features such as habit, leaf colour, leaf shape etc. but differs in having short, broad, abruptly acuminate stem leaves with recurved margins and dentate costa; paraphyllia with lateral papillae at the cell ends. However, *T. orientale* has erect or erect-spreading, ovate – lanceolate stem leaves with branched, filamentous, papillose paraphyllia. The species also resembles in most of the characters of *T. glaucinum* (Mitten) Bosch & Lacoste but differs in structure of costa and paraphyllia. In *T. glaucinum* costa ends below the tip and shows a tendency of bifurcation at tip and paraphyllia branched, lanceolate to filiform papillose. The Nagaland specimens show major resemblance in the characters of *T. recognitum* (Hedw.) Lindb. var. *delicatulum* (Hedw.) Warnst as described by earlier workers (Watanabe, 1972; Gangulee, 1978; Vohra, 1983) though some minor variations in the size of plants, leaves etc. which may be due to different ecological and geographical conditions.

ACKNOWLEDGEMENTS

Thanks are due to the Director, National Botanical Research Institute, Lucknow (India) for providing facilities and Council of Scientific and Industrial Research, New Delhi

for awarding Emeritus Scientist Scheme to Senior author (VN). Prof. S. K. Chaturvedi, Department of Botany, Nagaland University, Nagaland, is also acknowledged for providing necessary facilities during field trip.

LITERATURE CITED

- Bruch, P., W. P. Schimper and W. T. Gumbel.** 1836-1866. *Bryologia Europaea, Seu genera Muscorum Europaeorum monographiae illustrata* (Volume 6, fasc.65) cum corollaire et indice, Acced. Supplementum Schimper, Musci Europaei novi vel Bryologiae Supplementum Stuttgart.
- Crum, H. A.** 1999. *Bryophyte Flora of North America*. Provisional Publication.
- Gangulee, H. C.** 1978. *Mosses of Eastern India and Adjacent Regions*. Vol. III. Books and Allied (P) Ltd., Kolkata, India. pp. 1618-1649.
- Kuo, C.-M. and T.-Y. Chiang.** 1987. Index of Taiwan Mosses. *Taiwania* **32**: 119-207.
- Kürschner, H. and A. Erdağ.** 2005. Bryophytes of Turkey: An annotated reference list of the species with synonyms from the recent literature and an annotated list of Turkish Bryological Literature. *Turk. J. Bot.* **29**: 95-154.
- Lal, J.** 2005. A Checklist of Indian Mosses. Bishen Singh Mahendra Pal Singh, Dehra Dun, India. pp. 132-134.
- Noguchi, A., Z. Iwatsuki and T. Yamaguchi.** 1991. *Illustrated Moss Flora of Japan*. Vol. IV. The Hattori Botanical Laboratory, Japan.
- Norris, D. H. and T. Koponen.** 1985. Bryophyte flora of the Huon Peninsula, Papua New Guinea. VII. Trachypodaceae, Thuidiaceae and Meteoriaceae. *Acta bot. Fenn.* **131**: 1-51.
- Robinson, H.** 1968. Notes on the bryophytes from the Himalayas and Assam. *Bryologist* **72**: 82-97.
- Touw, A.** 1976. A taxonomic revision of *Thuidium*, *Pelekium*, and *Raiella* (Musci: Thuidiaceae) in Africa South of the Sahara. *Lindbergia* **3**: 135-195.
- Touw, A.** 2001. A review of the Thuidiaceae (Musci) and a realignment of taxa traditionally accommodated in *Thuidium sensu amplo* (*Thuidium* Schimp., *Thuidiopsis* (Broth.) M. Fleisch. and *Pelekium* Mitt.) including *Aequatoriella* gen. nov. and *Indothuidium* gen. nov. *J. Hattori Bot. Lab.* **90**: 167-209.
- Touw, A. and L. F.-V. D. Haak.** 1989. A revision of the Australasian Thuidiaceae (Musci), with notes on species from adjacent regions. *J. Hattori Bot. Lab.* **67**: 1-57.
- Vohra, J. N.** 1983. *Hypnobryales* suborder Leskeineae (Musci) of the Himalayas. *Rec. Bot. Surv. India* **23**: 103-146.
- Watanabe, R.** 1972. A revision of the family Thuidiaceae in Japan and adjacent area. *J. Hattori Bot. Lab.* **36**: 171-320.
- Watanabe, R.** 1991. Notes on the Thuidiaceae in Asia. *J. Hattori Bot. Lab.* **69**: 37-47.



印度蘚類植物的新紀錄 – 鈎葉羽蘚 (*Thuidium recognitum* (Hedw.) Lindb. var. *delicatulum* (Hedw.) Warnst)

Virendra Nath^(1*), Pooja Bansal⁽¹⁾ and A. K. Asthana⁽¹⁾

1. Bryology Laboratory, National Botanical Research Institute (Council of Scientific and Industrial Research, New Delhi, India) Lucknow – 226 001, India.

** Corresponding author. Tel.: 0522-2297837; Fax: 0522-2205836, 2205839; Email: drvirendranath2001@rediffmail.com*

(收稿日期：2011 年 4 月 7 日；接受日期：2011 年 5 月 23 日)

摘要：本文報導屬於蘚類羽蘚科的鈎葉羽蘚首次於印度被發現。這份標本採集於在那加蘭邦莫戈格瓊的迪庫河，與鈎葉羽蘚一樣莖葉為銳尖形，莖葉上具有頂端圓椎狀的乳突，其葉主脈為鋸齒狀延伸未達葉尖。該種為印度苔蘚誌的新增補，但僅侷限分布於喜馬拉雅的東部。

關鍵詞：東喜馬拉雅、印度、蘚類、新紀錄、羽蘚科、羽蘚屬。