

NOTES ON THE FLORA OF TAIWAN (20)-- *SCUTELLARIA* (LAMIACEAE) IN TAIWAN⁽¹⁾

Tsung-Hsin Hsieh⁽²⁾ and Tseng-Chieng Huang^(2,3)

(Manuscript received 1 December 1994; accepted 26 January 1995)

ABSTRACT: A brief revision of the species of *Scutellaria* (Lamiaceae) in Taiwan has been made. Five species are accepted in this paper. Three corolla types including the robust type, geniculate type, and curved type can be distinguished. Four types of nutlet coat ornamentation including grape-hook type, radiated umbrella-like type, rounded concentric type, and mammillate type are observed. SEM micro-graphs of pollen grains and nutlets, chromosome numbers, key to the species, species illustrations, distribution maps, and other relevant informations, are provided.

KEYWORDS: *Scutellaria*, Lamiaceae, revision, Taiwan.

INTRODUCTION

Scutellaria, with 425 species (Paton 1990), is a large genus of the Lamiaceae. It was first reported by A. Henry (1896) in Taiwan. Since then, several papers on *Scutellaria* of Taiwan have been published by Matsumura and Hayata (1906), Hayata (1911, 1919), Kudo (1929), Yamamoto (1934), Ohwi (1934, 1935), Sasaki (1936), Yamazaki (1969, 1992), Li (1977), Huang and Cheng (1978), and Ying (1991).

Although many papers were published, some questions are still unanswered. The specimen of Henry 2073 (identified as *Scutellaria indica* L.) collected from S. Cape (Southern Taiwan) is dubious, because no specimen of *S. indica* from Southern Taiwan has been found since Henry. *Scutellaria taiwanensis* C. Y. Wu was reported by C. Y. Wu in 1977 according to the specimen Kudo et Mori s. n. 1931 from Taiwan, but it is not documented. *Scutellaria tarokoensis* Yamazaki was distinguished from *S. tashiroi* Hayata by Yamazaki in 1992. The former has upcurved short pubescence on stem and small leaves (0.5-2 x 0.4-1.5 cm), and the latter has retrorsely short pubescence on stem and large leaves (2-3.5 x 1.5-2.5 cm). Based on the specimens in TAI, the leaves of *S. tashiroi* and *S. tarokoensis* are variable and highly overlapping. In our opinion, leaf size is not a good characteristic. According to these points, the genus *Scutellaria* needs to be clarified in Taiwan.

(1) This work was supported by the research grant to T. C. Huang from the National Science Council, R. O. China(NSC82-0211-B002-178).

(2) Department of Botany, National Taiwan University, Taipei, Taiwan, Republic of China.

(3) Corresponding author.

Such features, as floral morphology, pollen grains, and nutlets, are discussed. Distribution maps, descriptions and a key to the species are also given.

MATERIALS AND METHODS

Specimens examined in this study are listed with acronyms of the herbaria excluding TAI. Pollen grains prepared by the Erdtman method (1952) were studied with LM and SEM. Somatic chromosomes were studied in root tips which had been held in a solution of 0.002 M 8-hydroxyquinoline for 3-4 h at a temperature of 18-20 °C, and then fixed in 1:3 acetic ethanol over night, hydrolysed in pectinase and squashed in acetic orcein. The voucher specimens are listed in Table 1.

RESULTS

(a). Floral morphology

Based on the length of the corolla tubes and their geniculate features three types of floral morphology are recognized in Taiwan (Fig. 1; Table 1). The robust type such as in *Scutellaria tashiroi* has a longer floral tube (2.0-2.5 cm) which is a little curved at the base. The geniculate type found in *S. indica* and *S. playfari* has a medium length tube (1.5-2.0 cm long) which becomes abruptly vertical at the base. The curved type, in *S. barbata* and *S. taiwanensis*, has a shorter floral tube (0.9-1.5 cm) which is curved at the base but never at right angles.

(b). Pollen morphology

The pollen morphology of all Taiwan species is isopolar, tricolporate, subprolate to prolate, with rugular to reticulate tectate surface with microperforate ornamentation (Fig. 2; Table 2). The ornamentation of the grains in the genus is the same, but *Scutellaria taiwanensis* has the largest pollen grains (Fig. 3; Table 2) among the species..

(c). Nutlet morphology

It is easy to distinguish all the Taiwan *Scutellaria* species by their nutlets, especially by the ornamentation of the nutlet coat. The ornamentation of nutlet coats is composed chiefly of various arrangements of tuberculae, but the term "tuberculae" used in this paper may be interpreted differently from that of previous specialists who used light microscopes or magnifying glasses. Based on the grouping pattern of tuberculae and the shape of elevated tuberculae within the reticulae, four types of ornamentations are proposed: Type 1, grappling-hook type, has elevated tuberculae like grappling-hooks on the nutlet coat as in *Scutellaria tashiroi* (Fig. 4C; Table 2). Type 2, radiated umbrella-like

type, has elevated tuberculae radiating umbrella-like as in *Scutellaria barbata* (Fig. 4F) and *S. indica* nutlets (Fig. 4I), but the hilum of the former (Fig. 4E) is shorter than that of the latter (Fig. 4H). Type 3, rounded concentric type, has rounded concentric tuberculae as in *Scutellaria playfairi* nutlets (Fig. C). Type 4, mammillate type, has mammilae-like elevated tuberculae as in nutlets of *Scutellaria taiwanensis* (Fig. 5F). The grouping pattern of tuberculae in types 1 to 3 is reticulate whereas in type 4 it is non-reticulate.

(d). Chromosome

Chromosome numbers of Subgen. *Scutellaria* have been reported as $2n = 24\text{-}34$ (Paton, 1990). In this study, the chromosomes of all five species are $2n = 26$ (Fig. 6) and extremely small.




Fig. 1. Floral morphology of the species of *Scutellaria indica* L. (a), *S. playfari* Kudo (b), *S. tashiroi* Hay. (c), *S. barbata* D. Don. (d), and *S. taiwanensis* C. Y. Wu (e).

Table 1. Comparison of floral morphology of *Scutellaria* in Taiwan.

Character Taxon	Color	Size (cm)	Corolla form at base	Voucher specimen
<i>S. barbata</i>	purple	0.9--1.3	little curved	<i>Hsieh 901</i>
<i>S. indica</i>	purple, pink, or white	1.5--2.0	geniculate	<i>Hsieh 892</i>
<i>S. playfairi</i>	whitish purple	1.5--1.8	geniculate	<i>Hsieh 1146</i>
<i>S. taiwanensis</i>	white, throat with purple spot	1.0--1.5	curved	<i>Hsieh 1011</i>
<i>S. tashiroi</i>	dark purple	2.0--2.5	little curved	<i>Hsieh 1181, 1320</i>

Table 2. Comparison of pollen grains and seed morphology of *Scutellaria* in Taiwan.

Character Taxon	Size of pollen (μ m)	Exine pattern	Size of nutlet (mm)	Type of nutlet coat
<i>S. barbata</i>	20-24 x 12-16	reticulate	1.1 x 0.8	radiated umbrella-like type
<i>S. indica</i>	18-23 x 12-17	finely reticulate	1.3 x 1.0	radiated umbrella-like type
<i>S. playfairi</i>	16-20 x 10-15	loose reticulate to rugulate	1.0 x 0.8	rounded concentric tuberculae type
<i>S. taiwanensis</i>	23-30 x 16-20	irregular rugulate	1.4 x 1.2	mammillate type
<i>S. tashiroi</i>	18-22 x 14-17	loose reticulate to rugulate	0.9 x 0.8	grappling-hook type




Fig. 2. SEM micrographs of pollen grains of *Scutellaria playfarii* Kudo (A, B), *S. barbata* D. Don. (C, D), and *S. indica* L. (E, F). White scale bar = 3 μ m. Black scale bar = 1 μ m.




Fig. 3. SEM micrographs of pollen grains of *Scutellaria tashiroi* Hayata (A, B) and *S. taiwanensis* (C, D). White scale bar = 3 μ m. Black scale bar = 1 μ m.




Fig. 4. SEM micrographs of seeds of *Scutellaria tashiroi* Hayata (A, B, C), *S. barbata* D. Don. (D, E, F), and *S. indica* L. (G, H, I). White scale bar = 200 μ m. Black scale bar = 30 μ m.




Fig. 5. SEM micrographs of seeds of *Scutellaria playfairi* Kudo (A, B, C) and *S. taiwanensis* C. Y. Wu (D, E, F). White scale bar = 200 μ m. Black scale bars = 30 μ m.




Fig. 6. Mitotic metaphase chromosome of *Scutellaria barbata* D. Don. (A), *S. indica* L. (B), *S. playfairi* Kudo (C), *S. taiwanensis* C Y. Wu (D), and *S. tashiroi* Hayata (E). Scale bar = 4 μ m.

CONCLUSION

After our studies in floral morphology and of palynological and seed morphological features, we arrived at the following taxonomic treatment. Five species in Taiwan are recognized. A key to species preceding the description of each taxon is presented:

KEY TO THE SPECIES

1. Leaves narrowly ovate, glabrous on both surfaces..... (1) *S. barbata*
1. Leaves ovate to broadly ovate, pubescent at least on the veins of the lower surface
 2. Petioles short, about 2-5-(8) mm long
 3. Corolla tube geniculate on the anterior side at base, 1.5--1.8 cm long..... (3) *S. playfairi*
 3. Corolla tube slightly curved on the anterior side at base, 2.0-2.5 cm long..... (5) *S. tashiroi*
 2. Petioles long, about 10-20 mm long
 4. Leaves densely pubescent ; corolla tube 1.5 -2.0 cm long..... (2) *S. indica*
 4. Leaves glabrous or pubescent on the veins of the lower surface; corolla tube 1.0 -1.5 cm long..... (4) *S. taiwanensis*

1. ***Scutellaria barbata*** D. Don, Prod. Fl. Nepal. 09. 1825; Ohwi, Act. Phytotax. Geobot. 4: 32. 1935; Murata in Hara eds. Fl. E. Himal. 2nd Rep. 117. 1971; H. W. Li, Fl. Reip. Pop. Sinic. 65(2): 229. t. 49-5. 1977; Yamazaki in Journ. Jap. Bot. 67: 315-319. 1992. 向天盞 Fig. 7

Scutellaria rivularis Benth. in Wall., Cat. n. 214. 1829; Benth. in Wall., Pl. As. Rar. 1: 66. 1830; Matsum. et Hayata in Journ. Coll. Sci. Univ. Tokyo 22: 314. 1906; Kudo, Mem. Fac. Sci. Agr. Taihoku Univ. 2: 268. 1929; Huang et Cheng, Fl. Taiwan 4: 524. 1978.

A perennial herb, glabrous, up to 30 cm high; stems erect, square in x-section. Leaves sessile or lower ones with petioles up to 0.5 cm long; blades narrowly ovate to ovate, 1-3 x 0.5-1 cm, the base obtuse to truncate, the apex acute to obtuse, the margins distantly serrate on the lower leaves, entire on the upper ones, glabrous, both surfaces covered with glandular dots, the veinlets 2-3 paired. Flowers axillary in leafy bracts, opposite, forming a raceme; calyx tubular, bilobed, minutely hairy; corolla pinkish-purple, tubular, 9-13 mm long, bilabiate, covered with glandular hairs, the upper lip very broadly ovate, emarginate at apex, the lower lip broadly ovate, 3-lobed, the central lobe emarginate at apex. Stamens 4, didynamous. Pollen grains 3-colporate, prolate, isopolar, 20-24 x 12-16 um, tectum reticulate with microperforae. Nutlets orbicular, 1.1 x 0.8 mm, nutlet coat with umbrella-like elevated tuberculae as ornamentation; hilum lateral.

Flowering and fruiting season from October to the following May.

This species is mainly distributed in the northern area and rarely in the southern area.

Specimens examined:

Taipei Co.: Chou et al. 7633, Yamamoto s. n. Dec 1930; Chisingshan, Wu s. n. May 1934; Hsiaokotou, Tang 1362, Huang 214, Hsu s. n. Dec 1968, Hsieh 901; Kankou, Liu et Shen s. n. Sept 1933; Keelung, Suzuki 4486, Sasaki s. n. May 1925; Mucha, Peng 1464; Pali, Ho 191; Pinlin, Kuo 8119; Shihting, T. C. Huang 10372, Kuo 11561; Suigenchi, Shimizu 210, Suzuki s. n. Apr 1929; Tamsui, Hsu 4441, T. Y. Yang et K. C. Yang 4474; Wantang, Cheng 737; Wuchishan, Cheng 1478; ihoton, Kao 7621. **Ilan Co.:** Chilan, T.




Fig. 7. Latitudinal and altitudinal distribution of *Scutellaria barbata* D. Don.

C. Huang et al. 10725; Tali, *Chang* 2171; Shanshin, *Kao* 9656. **Taoyuan Co.:** Chungli, *Kuo* 6613; Jenmei, *Kuo* 6197; Palin, *Chiang* s. n. Apr 1990. **Miaoli Co.:** *T. C. Huang* 5277. **Taichung Co.:** Fongyuan, *Sasaki* s. n. Jan 1923; Pingtung, *Hsu* 9128; Tachia, *Kao* 9677. **Kaohsiung Co.:** Kaohsiung, *Playfair* s. n. May 1889 (K, Photo). **Pingtung Co.:** Pingtung, S. Z. *Yang* 1041, 1042.

2. **Scutellaria indica** L., Sp. Pl. 600. 1753; Matsum. et Hayata, Journ. Coll. Sci. Univ. Tokyo **22**: 313. 1906; Hayata, Icon. Pl. Formos. **6**: 58. 1916. et 8: 85. 1919; . W. Li, Fl. Reip. Pop. Sinic. **65**(2): 172. 1977. excl. syn. *S. tashiroi* Hayata; Huang et Cheng. Fl. Taiwan **4**: 521. 1978.

印度黃芩 Fig. 8

Scutellaria taiwanense Ying, Mem. Coll. Agr. NTU. **31**(1): 24. 1991. *syn. nov.*

Type: Taichung Co. Derchi, *Ying* s. n. Sept 1990. (NTUF!).

A perennial herb, up to 20 cm high, covered with septate hairs; stems erect or procumbent at base, often tufted, puberulent. Leaves chartaceous, petiolate; petioles puberulent, 1-2 cm long; blades broadly ovate to cordate, 1.5-4 x 1-3.5 cm, the base obtuse to truncate, the apex obtuse, the margins crenate, puberulent, veinlets 4-5 pairs. Inflorescence a terminal raceme, up to 8 cm long; leafy bracts small, obovate. puberulent and glandular-hairy, bearing a scutellum in front; corolla purple, tubular, 1.5-2 cm long, prominently geniculate at base, bilabiate. Pollen grains 3-colporate, subprolate, isopolar, 18-23 x 12-17 um, tectum finely reticulate with microperforae. Nutlets ellipsoidal, 1.3 x 1.0 mm, nutlet coat with umbrella-like elevated tuberculae ornamentation; hilum about one third the length of the seed.

Flowering and fruiting season from September to the following May.

This species is distributed in northern and central areas below 2,400 m elevation in Taiwan.

Specimens examined:

Taipei Co.: Fulon, *Kao* 10727; Hsintien, *Shimizu* 2074; Kanko, *Cheng* 480; Keelung, *Suzuki* 4711, *Sasaki* s. n. Mar. 1930; Kueishan, *Suzuki* s. n. Feb. 1928, *Chuang* 3161; Kuangingshan, *Kuo* 6078, *Chuang* 2155; *Mutsushan*, *Wang* 3016, *Shimitsu* 505; Neihu, *Kao* 4494; Peihsinchuang, *Ou et Kao* 9278; Pinlin, *T. C. Huang* 7963, *H. N. Yang* s. n. Nov. 1978, *Chou* 10; Shihting, *T. C. Huang* 10371; Suangchikou, *Kuo* 10866; Tatunshan, *T. C. Huang* 9073, *Suzuki* 10244; Wulai, *Kao* 7170, *Hsieh* 892, *Sasaki* s. n. Mar. 1922; Yangmingshan, *Kuo* 2903, *Chuang* 2155. **Ilan Co.:** Lotung, *Sasaki* s. n. Oct. 1928. **Taoyuan Co.:** Amuhyo, *Mori* 266; Palin, *Cheng* 649. **Hsinchu Co.:** Tachienshan, *H. N. Yang* 1598, Hsienchiaoshih, *Hsieh* 858. **Miaoli Co.:** Mt. Santou, *T. C. Huang* 8792; Sansa, *Shimada* 1127. **Taichung Co.:** Kukan, *T. C. Huang* 10325, 14549; Kukan to Chingshan, *T. Y. Yang* 1358, *J. C. Wang* 2927; Pahsienshan, *Suzuki* 5580; Shunkuan, *T. C. Huang* et *C. F. Hsieh* 7303; Heuhshan, *Ying* 5103(NTUF). **Nantou Co.:** Chitou, *T.C. Huang* 9; Kuangtaochi, *C. F. Hsieh* et *Wu* 1438; Meyfong, *Ou et Kao* 9366; Puli, *Seki* 105; Tangpu, *S. Z. Yang* 3766 (PAI); Tungan, *Cheng* 1390. **Hualien Co.:** Taroko, *Suzuki* 8894; Tiensiang, *Hsu* 492; Yenhai logging trail, *J. C. Wang* 7613.




Fig. 8. Latitudinal and altitudinal distribution of *Scutellaria indica* L.

3. **Scutellaria playfairi** Kudo, Mem. Fac. Sci. Agr. Taihoku Univ. **2**: 254. 1929; Yamazaki in Journ. Jap. Bot. 67: 316. 1992. p.p., excl. syn. *S. procumbens* Ohwi.

布烈氏黃芩 Fig. 9, 10.

Scutellaria luzonica auct. non Rolfe: Forb. et Hemsl. in Journ. Linn. Soc. Bot. **26**: 296. 1890; Henry, List Pl. Formos. 73. 1896; Matsum. et Hayata in Journ. Coll. Sci. Univ. Tokyo **22**: 314. 1906; Hayata in Journ. Coll. Sci. Univ. Tokyo **25**: 183. 1908; Sasaki, List Pl. Formos. 360. 1928.

Scutellaria tashiroi Hayata var. *playfairi* (Kudo) Yamazaki in Journ. Jap. Bot. **44**(6): 174. 1969. excl. syn. *S. procumbens* Ohwi.

Scutellaria luzonica Rolfe var. *playfairi* (Kudo) Yamamoto in Journ. Trop. Agr. **6**: 558. 1934.

Scutellaria playfairi auct. non Kudo: Li, Fl. Reip. Pop. Sinic. **65**(2): 146. 1977.

Scutellaria javanica Junch. var. *playfairi* (Kudo) Huang et Cheng in Fl. Taiwan **4**: 523. 1978. excl. Hualien. Co. specimens.

A perennial herb, up to 40 cm high; stems square in x-section, pubescent. Leaves coriaceous, petiolate; petioles pubescent, 2-5(-8) mm long; blades ovate, 1-1.5 x 1.5-1.8 cm, the base obtuse to round, the apex obtuse or round, the margins serrate, the upper surface glabrous, the veinlets sunken, the lower surface purplish, pubescent, glandular-dotted, the veinlets prominent beneath. Inflorescence in a terminal raceme, flowers opposite, 4-20 cm long, pedicels twisted, nearly in one plane, corolla purplish blue, tubular, 1.3 cm long, prominently geniculate at base, bilabiate. Pollen grains 3-colporate, prolate, 16-20 x 10-15 um, tectum loose reticulate to rugulate with microperforae. Nutlets orbicular, 1.0 x 0.8 mm, nutlet coat with rounded concentric elevated tuberculae ornamentation; hilum lateral.

Flowering and fruiting season from June to December.

S. playfairi and *S. tashiroi* are closely related in external morphology. They have been considered as different varieties in the same species, until Yamazaki (1992) segregated them. Our results support this latter opinion.

Specimens examined:

Chiayi Co.: Fanlu, Peng 13449 (HAST). **Tainan Co.:** Paiho, Kuoh 3945; Kuantzuling, Morimoto 486, A. Hsieh 9. **Kaohsiung Co.:** Chishan, Mori s. n. Nov. 1907; Fongshan, Moritami s. n. Nov. 1910; Liukuei, Tsai 26; Tengchih, T. C. Huang et al. 14434. **Pingtung Co.:** Hengchun, T. C. Huang 13351, Suzuki 6133, 4978; Machia, T. C. Huang et Kao 7357; Nanjenshan, Suzuki 5022, 6088, Matsuda s. n. Aug. 1915, Shihwen, Hsieh 1146; Shoka to Tsaopu, T. C. Huang et al. 14076; Tachienshihshan, Peng 7142 (HAST); Lilongshan, T. C. Huang et al. 16266; enroute from Wutai to Haucha, Lin 380 (HAST). **Taitung Co.:** Luyehchi, Yamamoto et Mori 12, 65; Hsinwulu, S. F. Huang 3200, 3698; Tawu, Shimizu 2789; Tawu to Tsusueipo, Hsu 3432.

4. **Scutellaria taiwanensis** C. Y. Wu, Fl. Reip. Pop. Sinic. **65**(2): 177, 580. t. 40, 15-21. 1977. Type : In silvis Arayawan, Kudo et Mori s. n. Apr. 1930 (Holotype PE!).

臺灣黃芩 Fig. 9, 11.

Scutellaria indica auct. non L. : Henry in List Pl. Formos. Henry 896.

Scutellaria lilungense Ying in Mem. Coll. Agr. NTU. **31**(1): 26-27. 1991. Type: Lilongshan, Pingtung Co.: Ying s. n. Nov. 18. 1990. (Holotype NTUF!). *syn. nov.*




Fig. 9. Latitudinal and altitudinal distribution of *Scutellaria playfairi* Kudo (*), *S. taiwanensis* C. Y. Wu (★), and *S. tashiroi* Hayata (●).




Fig. 10. Illustration of *Scutellaria playfairi* Kudo. 1. Habit; 2. Leaf; 3. Corolla; 4. Anther; 5. Pistil; 6. Fruits.




Fig. 11. Illustration of *Scutellaria taiwanensis* C. Y. Wu. 1. Habit; 2. Cross-section of stem; 3. Leaf; 4. Hairs of leaf; 5. Corolla; 6. Fruits.

A perennial herb, 10-25 cm high; stems erect, often tufted, pubescent. Leaves chartaceous, petiolate; petioles pubescent, 0.5-1.5 cm long; blades ovate to rhomboid-ovate, 1.5-3.0 x 1.0-2.0 cm, the base obtuse or truncate, the apex acuminate to obtuse, the margins crenate 2-4, glabrous, the lower surface purple, 3-4 nerved. Flowers 4-10, opposite, forming a terminal loose raceme, 2-4 cm long, bracts leafy ovate, small, entire and ciliate; corolla white with purplish spot in throat, 1-1.5 cm long, pubescent. Pollen grains 3-colporate, subprolate, isopolar, 23-30 x 16-20 um, tectum irregular rugulate with microperforae. Nutlets orbicular, 1.4 x 1.2 mm; nutlet coat with mammilae tuberculae ornamentation.

Flowering and fruiting season from September to the following April.

The specimen identified by A. Henry (*Henry 2073, K*) as *S. indica* L. (as listed in his List of Plant Formos. n. 802.), should be correctly referred to *S. taiwanensis*. The type specimen of *S. taiwanensis* C. Y. Wu collected by *Kudo et Mori s. n.* Apr. 1930 from Arayuhan, is located in the southern part of Taiwan, but precise locality is not known now. So, this species attributed to Mt. Ali by Li (1977) is a mistake.

Specimens examined:

Pingtung Co.: Lilongshan, Ying s. n. Nov. 1990 (Holotype of *S. lilungense* NTUF!), Lu 46 (PAI), S. Z. Yang 6621 (PAI), Hu 910, Huang et al. 16268; In silvis Arayuhan (southern Taiwan), *Kudo et Mori s. n.* Apr. 1930 (Holotype of *S. taiwanensis* PE!).

Taitung Co.: Chinshuiying, J. C. Wang et al. 7391 (TNU), S. Z. Yang et C. G. Lin 220720 (PAI), S. Z. Yang 25197, Hsieh 1011.

5. **Scutellaria tashiroi** Hayata, Icon. Pl. Formos. 8: 85. 1919; Yamazaki in Journ. Jap. Bot. 44(6): 175. 1969; et 67: 316. 1992. 田代氏黃芩 Fig. 9, 12.

Scutellaria procumbens Ohwi in Fedde Rep. Sp. Nov. 36: 52. 1934.

Scutellaria procumbens Ohwi var. *tomentosa* Ohwi in Act. Phyt. Geobot. 4: 33. 1935.

Scutellaria tashiroi Hatata var. *tomentosa* (Ohwi) Yamazaki in Journ. Jap. Bot. 44(6): 175. 1969.

Scutellaria playfairi auct. non Kudo.: H. W. Li, Fl. Reip. Pop. Sinic. 65(2): 146. 1977.

Scutellaria playfairi Kudo var. *procumbens* (Ohwi) Wu et Li. op. cit.

Scutellaria javanica Jungh. var. *luzonica* auct. non (Rolfe) Keng: Huang et Cheng in Fl. Taiwan 4:523. 1978. p. p.

Scutellaria tashiroi Hayata var. *haianshanensis* Yamazaki in Journ. Jap. Bot. 67: 316. 1992. *syn. nov.*

Scutellaria tarokoensis Yamazaki, Journ. Jap. Bot. 67: 316. 1992. *syn. nov.*

A suffruticose perennial herb, up to 100 cm long; stems slender, procumbent, tufted, puberulent. Leaves coriaceous, petiolate; petioles puberulent, 2-5(-8) mm long; blades ovate to triangular-ovate, (0.5-)2-3.5 x (0.4-)1.5-2.5 cm long, the apex obtuse, the base narrowly cordate, the margins crenate, the upper surface glabrous, the veinlets sunken, the lower surface puberulent or glabrous, covered with glandular dots. Inflorescence in axillary or terminal raceme, 3-8 cm long, flowers opposite, bracts cuneate-rhomboidal. Pollen grains 3-colporate, subprolate, isopolar, 18-22 x 14-17 um, tectum loose reticulate to rugulate with microperforae. Nutlets orbicular, 0.9 x 0.8 mm, nutlet coat with grappling-hook tuberculae ornamentation; hilum lateral.




Fig. 12. Illustration of *Scutellaria tashiroi* Hayata. 1. Habit; 2. Leaf; 3. Corolla; 4. Anther; 5. Pistil; 6. Fruits.

Flowering and fruiting season from July to December.

Although there is some variation in this disjunct species, an observation of the living plants and a study of a large number of herbarium specimens indicate that the varieties are conspecific with reference to the floral morphology, pollen grains, and seed morphology.

Specimens examined:

Hualien Co.: Taroko, Hosokawa 5113, Suzuki s. n. Nov. 1932, Suzuki et Hukuyama 16352, Keng et Kao 2586; Tienhsian, T. C. Huang 10169, Hsieh 1181; Chilai, Suzuki 1878, Sasaki s. n. Aug. 1929. Nakamura 4925, 4994; Luanshan, T. C. Huang 4248, Kuo et al. 6955, C. Hsu 3630; Chienching, Suzuki s. n. Aug. 1932; Takimi, Suzuki 1508; Yushan, Sasaki s. n. Aug. 1935(without further locality).
Taitung Co.: Luyehchi, Yamamoto et Mori 285; Kwanshan, S. Z. Yang 30372, 30323; **Orchid Island:** T. C. Huang et Kao 5370, 5439, C. F. Hsieh 1614; Hungtou to Tientzu, Chang 2272; Mt. Sentaku, Sata 1677, 1678, 1679; Tasenshan, Hosokawa s. n. July 1935; Tienchu, Hsieh 1320.

ACKNOWLEDGEMENT

We would like to thank Dr. T. Y. Yang who took several photos from specimens kept in Kew Herbarium and supplied us some important literatures. Gratitude is also due to the curators of the following herbaria, HAST, PAI, PE, NTUF, for allowing us examination of their specimens. H. P. Chuang, SEM Laboratory, Dept. of Botany, National Taiwan University, for taking SEM photographs, is also acknowledged.

LITERATURE CITED

- Hayata, A. 1911. Materials flora of Formos. Supplementary notes to the Enumerators plantarum Formosanum and flora montana Formosae, based on a study of the collections of the Botanical Survey of the government of Formosa, principally made at the herbarium of the Royal Botanical Gardens, Kew Journ. Coll. Sci. Univ. Tokyo **30**: 1- 471.
- Hayata, A. 1919. Icones of the plants of Formosa, and materials for Flora of the island, based on a study of the collections of the botanical survey of the Government of Formosa. **8**: 84-86
- Henry, A. 1896. A list of the plants from Formosa with some preliminary remarks on the geography, nature of the flora and Economic Botany of the island. Trans. Asiat. Soc. Jap. **24**. Suppl. 1-118.
- Huang, T. C. and W. T. Cheng 1978. Labiateae. In H. L. Li et al. eds., Flora of Taiwan. **4**: 39-529.
- Keng, H. 1969. Flora Malesianae Procursores XLVIII. A Revision of Malesian Labiateae. Guard. Bull. Sing. **24**: 13-180.

- Kudo, Y. 1929. Labiatarum Sino-Japonicarum Prodromus. Mem. Fac. Sci. Agr. Taihoku Univ. **2**(2): 37-332.
- Li, H. W. 1977. *Scutellaria*. In C. Y. Wu and H. W. Li, ed., Labiateae. Flora of Reip. Pop. Sinic. **65**(2): 124-248.
- Matsumura, J. and B. Hayata 1906. Enumeratato plantarum in Insula Formosae sponte crescentium huscusque rite congitarum adjectis descriptionibus et figuris specierum proregione novarum. Journ. Coll. Sci. Univ. Tokyo **22**: 305-320.
- Ohwi, J. 1934. Plantae Novae Japonicae. Fedde Rep. Sp. Nov. **36**: 52.
- Ohwi, J. 1935. Symbosae ad Floram Asiae Orientalis II. Act. Phyt. Geobot. **4**: 33.
- Paton, A. 1990. A global taxonomic investigation of *Scutellaria* (Labiatae). Kew Bull. **45**(3): 399-450.
- Yamazaki, T. 1969. Supplement of the flora of Ryukyu and Formosa. Journ. Jap. Bot. **44**(6): 174-175.
- Yamazaki, T. 1992. A revision of *Scutellaria* in Taiwan. Journ. Jap. Bot. **67**: 315-319.
- Ying, S. S. 1991. Miscellaneous notes on the flora of Taiwan XIV. Mem. Coll. Agr. NTU. **29**(1): 19-39.

臺灣植物誌之觀察(20)--臺灣產黃芩屬植物(唇形花科)⁽¹⁾

謝宗欣⁽²⁾、黃增泉^(2,3)

(收稿日期：1994年12月1日，接受日期：1995年1月26日)

摘要

本文使用外部形態、花粉、小堅果和地理分布等特徵，來探討本省產黃芩屬植物。結果可分為5種，即向天蓋、印度黃芩、布烈氏黃芩、臺灣黃芩和田代氏黃芩等。並指出小堅果的表皮微細構造為分種的良好特徵。本文並報導省產5種染色體數目皆為 $2n = 26$ 。

關鍵字：黃芩屬，唇形花科，分類訂正，臺灣。

1. 本文承國科會計劃補助(NSC82-0211-B002-178)。
2. 國立台灣大學植物學系。
3. 通信聯絡員。