

Oberonia segawae (Orchidaceae): A New Orchid Species in Taiwan

Tien-Chuan Hsu⁽¹⁾ and Shih-Wen Chung^(2,3)

(Manuscript received 17 October, 2007; accepted 22 January, 2008)

ABSTRACT: *Oberonia segawae*, a new orchid species found from central to south Taiwan, is described and illustrated. This species, which belongs to subgenus *Oberonia*, is characterized by its narrowly ensiform leaves up to 13 cm long and 5 mm wide, its whitish flowers, and its lip with indistinct lateral lobes and irregularly laciniate margins.

KEY WORDS: Orchidaceae, *Oberonia segawae*, new species, Taiwan, taxonomy.

INTRODUCTION

The orchid genus *Oberonia* Lindl. consists of about 300 species, distributed from South Africa, through Asia to Australia and the Pacific Islands. The plants of this genus are easily recognized by their distichous, laterally compressed leaves, and terminal racemes covered with numerous minute flowers. In Taiwan, five *Oberonia* species were listed in the latest overall revision (Su, 2000), and two small rhizomatic species were later transferred from *Hippeophyllum* (Ormerod, 2002). We recently discovered an *Oberonia* species with a lip shape quite distinct from the seven known species in Taiwan. After consulting the literature regarding *Oberonia* in neighboring regions, we conclude that the taxon constitutes a new species, which we describe and illustrate here.

Key to the species of *Oberonia* in Taiwan

- 1. Leaves jointed at base (subg. *Oberonia*)
 - 2. Leaves more than 1 cm wide; lip distinctly 3-lobed from base *O. gigantea*
 - 2. Leaves less than 0.8 cm wide; lip not distinctly 3-lobed from base.
 - 3. Stem 3-6 cm long; lip margin entire *O. caulescens*
 - 3. Stem less than 3 cm long; lip margin irregularly laciniate *O. segawae*
- 1. Leaves not jointed at base (subg. *Menophyllum*)
 - 4. Plants arising separately from distinct rhizome.
 - 5. Rhizomes loosely scaled; rachis not dilated; lip not 3-lobed from base *O. pumilum*
 - 5. Rhizomes densely scaled; rachis distinctly dilated; lip 3-lobed from base *O. seidenfadenii*
 - 4. Plants tufted, rhizome indistinct.
 - 6. Inflorescence shorter or as long as leaves; lateral lobes of lip extending upward, slightly surrounding column; petals distinctly erose *O. rosea segawae*

- 6. Inflorescence much longer than leaves; lateral lobes of lip extending apically, not surrounding column; petals entire or indistinctly erose.
- 7. Apical leaves 3-5 cm long, often much longer than central leaves; central lobe of lip rectangular *O. arisanensis*
- 7. Apical leaves 1-3 cm long, slightly longer than central leaves; central lobe of lip orbicular or squarish *O. japonica*

TAXONOMIC TREATMENTS

Oberonia segawae T.-C. Hsu & S.-W. Chung, *sp. nov.*
— TYPE: TAIWAN. Pingtung County, Chaozhoujun, Aug 1937, *K. Segawa s. n.* (holotype, TAIF 117872), here designated.

齒唇莖白蘭 Figs. 1-3

Species *O. gigantea* Fukuyama affinis, sed planta et floris minore, floris albidus, labellum lobi laterals indistinctus, marginis dentatum.

Perennial epiphyte. Plant often tufted and pendulous. Stems short, concealed by leaf sheaths, less than 3 cm long. Leaf 5-8, distichous, laterally compressed, narrowly ensiform, fleshy, variable in size, the largest ones 6-13 cm long, 0.3-0.5 mm wide, base jointed, apex sub-falcate. Inflorescence 5-10 cm long; peduncle short, 1.5-3 cm long, terete, with many small bracts; flowers numerous, arranged in whorls, forming a cylinder ca. 6 mm in diam. Bracts ovate-lanceolate, 2-2.5 mm long, 1-1.3 mm wide. Pedicel and ovary 1-2 mm long, finely ridged. Flowers minute, whitish or sometimes tinged with orange, ca. 1.5 mm in diam. Sepals spreading, subequal, entire, ovate-triangular, 0.7-0.8 mm long, 0.4-0.5 mm wide. Petals spreading, subentire, elliptic-oblong, 0.7-0.8 mm long, ca. 0.2 mm wide. Lip spreading, more or less perpendicular to ovary, slightly concave, ovate, nearly unlobed, base truncate, apex retuse, margin irregularly laciniate; the laciniate lobulets progressively deeper towards the apex. Column short, globular, ca. 0.3 mm

1. Division of Forestry Biology, Taiwan Forestry Research Institute, 53, Nanhai Road, Taipei 100, Taiwan.
 2. Department of Life Science, National Taiwan University, 1, Roosevelt Road, Sec. 4, Taipei 106, Taiwan.
 3. Corresponding author. Tel: 886-2-23039978; Email: biflora@gmail.com




Fig. 1. *Oberonia segawae* T. C. Hsu & S. W. Chung. A: Habit. B: Flower with ovary and bract, lateral view. C: Flower, front view. D: Dorsal sepal. E: Petal. F: Lateral sepal. G-H: Lips in varied shapes, cut from flowers in the same inflorescence.

long. Pollinia 4, in 2 pairs. Capsules ellipsoid, ca. 2 mm long.

Additional specimens examined: TAIWAN. Nantou County: Xingaojun, 25 Jun 1934, *K. Segawa s. n.* (TAIF); Ruan-an, ca. 1400 m, 27 Aug 2006, *T. -C. Hsu 567* (TAIF); Hsitou, 31 Jan 1986, *H. -J. Su 6953* (HAST). Chiayi County: Chiehtung Trail, ca. 1000 m, 15 Dec 2004, *P. -F. Lu 8854* (TAIF). Kaohsiung County: Zhonzhiguan, 3 May 1936, *K. Segawa s. n.* (TAIF); Sanhexi, 21 Jan 1934, *K. Segawa s. n.* (TAIF).

Distribution: Endemic to Taiwan. Presently known at elevations of 1000 to 2000 m a.s.l. in the

central and southern mountainous regions. Plants are usually found in moist forests along streams, growing as an epiphyte on branches of broadleaf trees or on vines.

Phenology: Flowers observed in August; fruits matured in September to October.

Etymology: The specific epithet is derived from the collector of the type specimen, Kokichi Segawa, a prominent plant collector and ethnologist.




Fig. 2. *Oberonia segawae* T. C. Hsu & S. W. Chung. A: Habitat and habits. B: Habit. C. Part of inflorescence with flowers.

Notes: This new species is assigned to subgenus *Oberonia* base on its jointed leaves (Seidendaden and Wood, 1992; Chen et al., 1999). In Taiwan, it can be easily distinguished from the two known species in this subgenus, *O. caulescens* Lindl. and *O. gigantean* Fukuyama (Lin, 1987; Chen et al., 1999; Su, 2000),

by its narrowly ensiform leaves to 13 cm long and 0.5 cm wide, and obscurely lobed lip which has an irregularly laciniate margin. Although specimens referable to this taxon have been collected several times since 1934, they were either determined as an unknown species or misidentified as *O. caulescens*.




Fig. 3. Distribution of *Oberonia segawae* (solid circles) in Taiwan.

There are certain Asian taxa, such as *O. mucronata* (D. Don) Ormerod & Seidenf., *O. denticulata* Wight (= *O. iridifolia* Lindl.), *O. gammiei* King & Pantl. and *O. dissitiflora* Ridl., which are characterized by jointed leaves, an obscurely lobed lip and an erose lip margin. However, after checking descriptions and illustrations regarding these species (Valmayor, 1984a, 1984b; Ansari and Balakrishnan, 1990; Seidendaden and Wood, 1992; Chen et al., 1999; Comber, 2001; Pearce and Cribb, 2002; Misra, 2004), none were found to adequately fit the material collected from Taiwan. We therefore regard it as a new species. Further systematic research on related species is needed, since the tiny flowers are problematic for comparative morphological studies.

The high fruit set ratio and rapidly expanding ovaries suggest the possibility of self-pollination in this species.

ACKNOWLEDGEMENTS

We are thanks to Mr. Sheng-Kun Yu and Ms. Pi-Fong Lu for their assistance in field works, and Dr. Tsan-Piao Lin, chairman/professor of Institute of Plant Biology, National Taiwan University, for kindly offering valuable information. We are indebted to Dr. Stephan Gale, The Kochi Prefectural Makino Botanical Garden, for reviewing English writing.

LITERATURE CITED

- Ansari, R. and N. P. Balakrishnan. 1990. A revision of the Indian species of *Oberonia* (Orchidaceae). *Orchid Monographs*, **4**: 82. Leiden University, The Netherlands.
- Chen X.-Q., J.-H. Ji, K.-Y. Lang and Y.-B. Lo. 1999. *Flora Reipublicae Popularis Sinicae*, **18**: 123-150. Science Press, Beijing, China.
- Comber, J. 2001. *Orchids of Sumatra*. Royal Botanic Gardens, Kew, UK. pp. 185-204.
- Lin, T.-P. 1987. *Native Orchids of Taiwan*, **3**: 177-196. Southern Materials Center, Taipei, Taiwan.
- Misra, S. 2004. *Orchids of Orissa*. Gajendra Singh Gahlot at Shive Offset Press, Dehra Dun, India. pp. 329-345.
- Pearce, N. R. and P. J. Cribb. 2002. *The Orchids of Bhutan*. Royal Botanic Gardens Edinburgh, Edinburgh, UK. pp. 119-120.
- Seidenfaden, G. and J. J. Wood. 1992. *The Orchids of Peninsular Malaysia and Singapore*. pp. 342-346. Olsen & Olsen, Fredensborg, Denmark. pp. 241-248.
- Su, H.-J. 2000. In: Huang, T.-C. et al. (eds.), *Orchidaceae. Flora of Taiwan*, 2nd ed. **5**: 981-985. Editorial Committee, Dept. Bot., NTU, Taipei, Taiwan.
- Valmayor, H. L. 1984a. *Orchidiana Philippiniana*, **1**: 253-257. Eugenio Lopez Foundation, Manila, Philippine.
- Valmayor, H. L. 1984b. *Orchidiana Philippiniana*, **2**: 117. Eugenio Lopez Foundation, Manila, Philippine.

臺灣新種蘭科植物：齒唇莪白蘭

許天銓⁽¹⁾、鐘詩文^(2,3)

(收稿日期：2007年10月17日；接受日期：2008年1月22日)

摘 要

本文描述產於臺灣中南部的蘭科新種：齒唇莪白蘭 *Oberonia segawae* T. C. Hsu & S. W. Chung。本種隸屬於 *Oberonia* 亞屬，且具有下列特徵：葉狹長劍形，長可達 12 公分，寬達 5 公釐；花近白色，唇瓣側裂片不顯著且邊緣呈現撕裂狀。

關鍵詞：蘭科、莪白蘭屬、齒唇莪白蘭、新種、臺灣、分類學。

1. 行政院農業委員會林業試驗所生物組，100 台北市南海路 53 號，臺灣。
2. 國立臺灣大學生態學與演化生物學研究所，106 台北市羅斯福路 4 段 1 號，臺灣。
3. 通信作者。Tel: 886-2-23039978; Email: biflora@gmail.com