



NOTE

Sikkim Himalayas – The Megadiversity Centre of *Pedicularis* L. in IndiaArti Garg^(1*) and Tariq Husain⁽²⁾

1. Central National Herbarium, Botanical Survey of India, P.O. Indian Botanic Gardens, Howrah 711 103, India. Email: kad_arti396@yahoo.com

2. Taxonomy and Biodiversity Division, National Botanical Research Institute, Rana Pratap Marg, Lucknow – 226 001 (U.P.), India. Email: Hustar_2000@yahoo.co.uk

* Corresponding author.

(Manuscript received 12 August 2009; accepted 16 November 2009)

ABSTRACT: Investigations on the Himalayan genus *Pedicularis* L. revealed rich concentration and enormous diversity of species in the Sikkim Himalayas. Out of the 83 Indian species of *Pedicularis*, 49 species (ca. 59%) are found in the Sikkim Himalayas and 34 species (ca. 41%) are endemic. The Sikkim Himalayas may therefore be designated as the ‘megadiversity centre’ of *Pedicularis* L. in India.

KEY WORDS: *Pedicularis*, Scrophulariaceae, Sikkim Himalayas, Endemic, Megadiversity centre.

INTRODUCTION

The Himalayan genus *Pedicularis* L. consists of eighty three species in India of which, 81 species are distributed in the Himalayan range, stretching from Ladakh in the west, through the Sikkim Himalayas, reaching up to Arunachal Pradesh in the east. Two species, namely *Pedicularis zeylanica* Benth. and *P. perrottetii* Benth., are confined to the Nilgiri Hills of south India. The genus *Pedicularis* L. was first described by Linnaeus in 1737 and subsequently by many workers viz. Bunge (1846), Maximowicz (1882), Pennell (1943) and Prain (1889, 1890) and was recently revised (Husain and Garg, 2007). The generic name *Pedicularis* derives its origin from the Latin ‘Pediculus’ meaning louse, as its decoction was used against lice on domestic animals in some areas of central Europe. Hence, the members of this genus are commonly known as louse-worts. Many species exhibit such natural variations and extreme adaptations in their floral dynamics which indicate their adaptive response to survive and perpetuate in the drastic climatic conditions native to them (Garg and Husain, 2003). (Fig. 1, Table 1)

The genus *Pedicularis* on the whole is restricted to the temperate, sub-alpine and alpine-montane zones, but a few species such as *Pedicularis bifida*, *P. punctata*, *P. hoffmeisteri* descend to 2000 meters. Some species (*P. lachnoglossa* Hook.f., *P. bicornuta* Kl. ex Kl. and Garcke, *P. hoffmeisteri* Kl. ex Kl. and Garcke and *Pedicularis rhinanthoides* Schrenk ex Fisch. & C.A. Mey.) often reach the highest altitudes of above 5000 meters, the maximum height reached by flowering plants. However, the majority of the species occur at altitudes between 2500-4500 meters (Li, 1951) and

rarely one or two species find their way towards the lowest extreme of 1000 meters (*Pedicularis khasiana* (Hook. f.) Pennell, *Pedicularis bifida* (Buch.- Ham. ex D. Don) Pennell).

Tsoong (1955a, b) reported the Sikang-Yunnan and the Sikkim regions to posses the highest concentration of *Pedicularis* species in the world. Yang Han-bi et al. (1998) concluded SW China to be the main center of diversity for *Pedicularis* species with about 352 species while few species occupy Bhutan (Mill, 2001). In India however, the maximum species concentration is found in the eastern Himalaya, especially in Sikkim extending in parts of China which projects in-between Sikkim and Bhutan. The number of species gradually decrease westwards. The species *P. punctata* Decne occupies most areas of the Kullu district extending from Kothi to Rohtang pass and further into the Lahaul and Spiti valley, whereas *Pedicularis bifida* (Buch. Ham. ex D. Don) Pennell occupies the entire Uttarakhand Himalayan regions. (Figs. 2 & 3)

The Sikkim Himalayas enjoy a position intermediate to the extreme climatic conditions of the eastern Himalayas on one hand and the western on the other. The eastern regions are endowed with uniformly moist stormy weather with frequent heavy rains, whereas the western zones are comparatively dry stormy and windy with ice-cold winds and frequent optimum rainfall. The magnitude of physical distance between the cold deserts of Ladakh and Lahaul-Spiti valley of western Himalaya and the Sikkim valley and the eastern Himalaya with a wide variety of ecological gradients induces variations among different species. The ideal location of the Sikkim Himalayas validate itself for species concentration and sustainence.



Fig. 1. Flower variation in species of *Pedicularis* occurring in Sikkim Himalayas. A: *P. megalantha* D. Don var. *megalantha*. B: *P. longiflora* Rudolf var. *tubiformis* (Klotzch) Tsoong. C: *P. roylei* Maxim. D: *P. siphonantha* D. Don. E: *P. rhinanthoides* Schrenk. ex Fisch. & Mey. subsp. *revoluta* Pennell. F: *P. porrecta* Wall. ex Benth.

MATERIAL AND METHODS

Field trips were made out to the Sikkim Himalayas during different flowering seasons to record the natural habitat, species occurrence and variations, ecological conditions, and microhabitat associations. All the collections were brought to laboratory, studied for the morphological variations among them and identified. Observations were recorded for plant morphology as related to diagnostic characters and morphological adaptations. The voucher specimens were deposited in

the Herbarium (LWG) of the National Botanical Research Institute, Lucknow, where the studies were undertaken.

RESULTS AND DISCUSSION

The natural abode of the *Pedicularis* species are regions which are covered with snow for most of the year. As snow starts melting at the onset of July the sprouting of underlying dormant rhizomes begin. Flowering and fruiting commences within a short time and the entire life cycle is completed in a short span of two months, after which the

**Table 1.** *Pedicularis* species occurring in the Sikkim Himalayas.

S. No.	Taxa	Distribution in India	Voucher specimen(s)
01*	<i>P. albiflora</i> (Hook.f.) Prain	Sikkim	Sikkim, Kankola, 13,000 ft., 22 Aug. 1879, J.D. Hooker s.n. (holo.: CAL!)
02*	<i>P. bella</i> Hook.f.	Sikkim	Sikkim, Darkia-la, 5660m, Aug., 1892, G.A. Gammie s.n. (DD).
03	<i>P. bifida</i> (Buch.-Ham. ex D. Don) Pennell	H. P., Uttaranchal, Sikkim, Assam, Meghalaya, Manipur	Uttaranchal, Kumaon, Verinag, <i>Hira Lal</i> 1514 (LWG); Kumaon, 1333 m, 21.8.1904, N. Gill 186 (LWG).
04*	<i>P. chumbica</i> Prain	Sikkim	Sikkim, Tanka-la, 5000 m, Aug., 1892, G.A. Gammie s.n. (DD); Seratong, 4500 m, 15.7.1910, W.W. Smith 3474 (CAL).
05	<i>P. clarkii</i> Hook. f.	Sikkim, Nagaland	Sikkim, Changu, 4000 m, 26.10.1910, <i>Ribu & Rhomoo</i> 4371 (CAL, two specimens); Nagaland, Naga hills, June 1936, <i>N.L. Bor</i> 20758 (ASSAM).
06	<i>P. collata</i> Prain	Sikkim, Meghalaya, Manipur	Sikkim, Jongri, 4000 m, Oct. 1898, s.l. 11974 (CAL); Lanepokui, 4500 m, 12.8.1913, <i>G.H. Cave</i> 905 (CAL).
07*	<i>P. cooperi</i> Tsoong	Sikkim	Sikkim, Chola Pass, 16 Sept. 1913, R.E. Cooper 884 (holo.: Herb. Edinb.-n.v.).
08*	<i>P. confertiflora</i> Prain	Sikkim	Sikkim, Tanqa, 4300 m, 15.8.1909, <i>Smith & Cave</i> 2550 (CAL)
09	<i>P. curvipes</i> Hook. f.	Sikkim, Nagaland, Manipur	Sikkim, Tumbok, 10,000 ft, 9.10.1870, C.B. Clarke s.n. (CAL, fragmentary isotype in packet, mounted on sheet also bearing <i>Clarke</i> 41334B from Nagaland).
10*	<i>P. daltoni</i> Prain	Sikkim	Sikkim, Jankrata, 5340 m, August 1892, G.A. Gammie s.n. (BM).
11*	<i>P. denudata</i> Hook.f.	Sikkim	Sikkim Himalaya, Lachen Valley, 12-13,000 ft, J.D. Hooker numbered 'Pedicularis no. 11'-n.v.
12*	<i>P. diffusa</i> Prain	Sikkim	Sikkim, above Toong, 3340 m, 18.9.1913, R.E. Cooper 905 (BM); Zenum valley, 3330 m, 10.7.1909, s.l. 1147 (CAL)
13*	<i>P. elwesii</i> Hook.f.	Sikkim	Sikkim, Lachung valley, Aug., 1892, G.A. Gammie s.n. (DD).
14*	<i>P. excelsa</i> Hook.f.	Sikkim	Sikkim, Yeumtong, 12, 000 ft, Sept. 1849, J.D. Hooker & Thomson <i>Pedicularis</i> no.27 (holo.: K-n.v.; iso.: BM!).
15*	<i>P. exigua</i> Li	Sikkim	Sikkim, Thangu, Sept. 1903, Prain s.n. (holo.: Univ. of California-n.v.; iso.: PH-n.v.).
16	<i>P. flagellaris</i> Benth.	Sikkim, Meghalaya	Meghalaya, K.&J. Hills, Laitlyngkot, 11.8.1938, R.N. De 16851 (ASSAM).
17*	<i>P. flexuosa</i> Hook.f.	Sikkim	Sikkim, Changu, 4000 m, 2.07.1913, R.E. Cooper 126 (BM); Kapup, 4340 m, 7.7.1913, R.E. Cooper 173 (BM)
18	<i>P. furfuracea</i> Wall.	West Bengal, Sikkim	Sikkim, June 1887, King s.n. (BM); Phusum, 3330 m, 25.6.1945, <i>Bor & Kirat Ram</i> 20586 (DD)
19*	<i>P. gammieana</i> Prain	Sikkim	Type: Sikkim, 'Eastern Himalaya, Lang-mang-nang-zo, 10,000ft, Pantling' (iso.: K-n.v.).
20	<i>P. gibbera</i> Prain	Sikkim, Arunachal Pradesh	Sikkim, Sept., 1887, Dr. King's Collector s.n. (DD); Sukki-moral, W. Tangui, Aug., 1887, Dr. King's collector s.n. (DD, CAL).
21*	<i>P. globifera</i> Hook.f.	Sikkim	Type: Sikkim Himalaya, Kongra Lama, 14-15,000ft, J.D. Hooker numbered 'Pedicularis no. 10'-n.v.
22a	<i>P. gracilis</i> Wall. subsp. <i>gracilis</i> var. <i>gracilis</i>	Kashmir to Sikkim, & Assam	Jammu & Kashmir, Bhadrawaha-Sevj, 2800 m, Aug. 1986, J.N. Vohra & B.D. Naithani 83032 (BSD). Himachal Pradesh, Manali, 2800 m, 13.8.1941, <i>N.L. Bor</i> 15326 (DD); Arunachal Pradesh, Bomdi la-Saleri, 2000 m, 19.10.1955, Rankin & Pretzlik 056 (BM).
22b	var. <i>stricta</i> (Wall. ex Prain) Husain & Garg	Kashmir to Sikkim	Uttaranchal, Kumaon, way to Ralam Glacier, 4000-4500 m, 15.9.1969, Pant & Naithani 39686 (BSD). Sikkim, 2000 m, R.H. Beddome 5752 (BM).
22c	Subsp. <i>brunoniana</i> (Wall. ex Pennell) Husain & Garg var. <i>brunoniana</i>	Himachal Pradesh to Sikkim &, Nagaland	Himachal Pradesh, Moralkanda, 3000 m, 29.9.1970, P.C. Sharma & D.P. Badola 880 (LWG); Sikkim, Tangloo, 2700-3300 m, 3.8.1862, T. Anderson 992 (DD)
22d	var. <i>macrocarpa</i> (Prain) Tsoong	Kashmir to Sikkim	Jammu & Kashmir, near Bhairon temple, 31.8.1989, B.P. Uniyal 77404 (BSD); Uttaranchal, Kumaon, Loharkhet, 2000-3220 m, 20.9.1957, T.A. Rao 4244 (BSD), Changu, 4333 m, 9.9.1913, R.E. Cooper 847 (BM).
23*	<i>P. garckeana</i> Prain ex Maxim.	Sikkim	Specimens from Chumbi and Phari are housed in DD (details not available)
24*	<i>P. instar</i> Prain ex Maxim.	Sikkim	Sikkim, Lachung valley, 5000 m, 15.8.1892, G.A. Gammie 789 (CAL).
25*	<i>P. integrifolia</i> Hook.f.	Sikkim	Type: Sikkim Himalaya, Tungu, 12-14,000 ft, J.D. Hooker numbered 'Pedicularis no. 8'-n.v.
26*	<i>P. kingii</i> Prain ex Maxim.	Sikkim	Sikkim, Poshing La, 4000 m, 21.7.1938, <i>Kingdon-Ward</i> 13952 (BM).
27*	<i>P. lachnoglossa</i> Hook.f.	Sikkim	Type: Sikkim, Lachen valley, 14,000 ft, 15.7.1849, J.D. Hooker numbered 'Pedicularis no. 32' (lecto.: K-n.v.-specimen with handwritten ticket; isolecto.: BM-n.v.).
28	<i>P. longiflora</i> Rudolf var. <i>tubiformis</i> (Klotzch) Tsoong	Jammu & Kashmir, Himachal Pradesh, Uttaranchal, Sikkim, Arunachal Pradesh, Assam	Jammu & Kashmir, Ladakh: above Tsakzham Tso, 5830 m, 23.7.1931, Walter Koelz 2415c (DD). Uttaranchal, Kumaon, Gorji valley, Parbhu, 12.8.1900, <i>Inayat</i> 24799 (DD); North Sikkim: Thangu to Lachen, 3200 m, 6.10.1997, P. Singh & S.S. Dash 20065 (BSHC).

**Table 1. Continued.**

S. No.	Taxa	Distribution in India	Voucher specimen(s)
29	<i>P. megalantha</i> D. Don var. <i>megalantha</i>	Sikkim, Uttarakhand	Uttarakhand, Garhwal, Tehri Garhwal, Rhudughera, 3670-4000 m, 19.7.1883, J.F.Duthie 237 (DD). Sikkim, Changu & Kapup, 4000-4333 m, 19.8.1913, R.E. Cooper 551 (BM).
30*	<i>P. megalantha</i> D. Don var. <i>pauciflora</i> Prain ex Maxim.	Sikkim	Type: Bloktan (Bhutan), near Tashi (Tagi), July, 1887, Dr. King's collector s.n. (holo.: CAL!; iso.: DD!). There is one more specimen of Dr. King's collector from Chumbi in CAL, of var. <i>pauciflora</i> collected in Oct. 1884. Sikkim, Kapup, 4340 m, 8.7.1913, R.E. Cooper 176 (BM); Patung lu, 17.7.1877, G. King 4395 (DD)
31*	<i>P. microcalyx</i> Hook.f.	Sikkim	Jammu & Kashmir, Ladakh, Shyok, 4200m, 29.8.1975, M.V. Vishwanathan 55042(BSD). Uttarakhand, Kumaon, Byans, Palang Gad, 3330-3660 m, 19.7.1886, J.F. Duthie 5844 (DD); Sikkim, 3300-4600 m, Hooker s.n. (BM).
32	<i>P. mollis</i> Wall	Jammu & Kashmir, Himachal Pradesh, Uttarakhand, Sikkim	Jammu & Kashmir, Ladakh, Shyok, 4200m, 29.8.1975, M.V. Vishwanathan 55042(BSD). Uttarakhand, Kumaon, Byans, Palang Gad, 3330-3660 m, 19.7.1886, J.F. Duthie 5844 (DD); Sikkim, 3300-4600 m, Hooker s.n. (BM).
33*	<i>P. nepalensis</i> Prain	Sikkim	Sikkim, Choon-goo, Sept. 1882, Dr. King's Collector s.n. (CAL).
34*	<i>P. odontophora</i> Prain	Sikkim	Sikkim, at Na-tong, 24.8.1878, Dungboo s.n. (DD, CAL).
35*	<i>P. oederi</i> Vahl subsp. <i>brachiophylla</i> (Pennell) Tsoong	Sikkim	Type: Sikkim, 16000 ft., 22.7.1909, Smith & Cave 1714 (holo.: PH-n.v., iso.: CAL!).
36*	<i>P. pantlingii</i> Prain	Sikkim	Sikkim, Ling-too, 4000m, 2.9.1882, King's collector s.n. (BM).
37*	<i>P. polygaloides</i> Hook.f.	Sikkim	Sikkim, Jang-ka-la, 14.8.1882, G. King's Collector s.n. (DD).
38	<i>P. porrecta</i> Wall.	Kashmir, Himachal Pradesh, Uttarakhand, Sikkim	Uttarakhand, Kumaon, Ralam valley, 16.8.1900, Inayat 24786 (DD); Ralam valley, 4500-5000 m, 24.8.1884, J.F. Duthie 3226 (DD)
39*	<i>P. Prainiana</i> Maxim.	Sikkim	Type: China, Chumbi, Lu-ma-poo, 20.7.1884, Dr. King's collector 509 (CAL!).
40*	<i>P. regelianae</i> Prain	Sikkim	Type: Sikkim, Jongri, 12,000 ft., Sept. 1887, Dr. King's Collector s.n. (lecto.: CAL!; isolecto.: BM!). Other original syntypes: Jongri, Tha-moo-tsa, 12000 ft., Aug. 1887, Dr. King's Collector s.n. (CAL!, BM!).
41*	<i>P. rhinanthoides</i> Schreb ex Fisch. & C.A.Mey. subsp. <i>revolute</i> Pennell	Sikkim	Type: Sikkim, Nlonok, Nak Chu, 5000-5500 m, 3 Aug. 1909, Smith & Cave 1945 (holo.: PH-n.v.; iso.: CAL!).
42*	<i>P. robusta</i> Hook.f.	Sikkim	Sikkim, Samdong, 5600 m, 17.8.1892, G.A.Gammie 873 (CAL).
43	<i>P. roylei</i> Maxim.	Jammu & Kashmir, Himachal Pradesh, Uttarakhand, Sikkim	Jammu & Kashmir, Chovsar, 17.7.1899, Inayat s.n. (DD); Sikkim, North Sikkim, Yum Thang (Hot water spring area), 19.7.1987, D.C.S. Raju & S. Singh 7733 (BSHC);
44*	<i>P. schizorrhyncha</i> Prain*	Sikkim	Sikkim, Kankerteng, 4300 m, 09.1888, Dr. King's Collector s.n. (BM, DD).
45*	<i>P. scullyana</i> Prain ex Maxim.*	Sikkim	Sikkim, Bijan, Dact Lefal, 3670 m, 09.1888, Dr. King's collector s.n. (BM, DD); Damkerka, Lefal forest, 4000m, 09.1888, Dr. King's collector s.n. (BM).
46	<i>P. siphonantha</i> D. Don - Fig. 1D	Jammu & Kashmir, Himachal Pradesh, Uttarakhand, Sikkim, Arunachal Pradesh, Assam	Jammu & Kashmir, Gulmarg: 3333 – 3700 m, 8.1922, J.H. Barbour s.n. (BM); Uttarakhand, Garhwal, Pangu, Stociczka 3 (BM).
47*	<i>P. sikkimensis</i> Bonati, W.W. Smith	Sikkim	Sikkim, Zeumn valley, 3900 m, 11.7.1909, Smith & Cave 1213 (CAL).
48*	<i>P. tenuicaulis</i> Prain	Sikkim	Sikkim, Guatung, Aug. 1891, Dr. Cannivas s.n. (CAL).
49	<i>P. trichoglossa</i> Hook.f.	Sikkim, Uttarakhand	Uttarakhand, Kumaon, Dhauli valley near Raina, 4000-4330m, 3.9.1884, J.F. Duthie 3222 (BM, DD); Sikkim, 4000-5000m, J.D. Hooker 29 (BM).

Note: Asterisk (*) indicate the species endemic to Sikkim

area is again covered by a thick blanket of snow towards the end of September. Boggie soil conditions are most suitable for the growth of *Pedicularis* but many species frequently occur in streams and nullahs.

A high degree of microhabitat-specific character was observed within the different species of *Pedicularis* with 4-5 plant species, predominantly of Legumes and some Graminae, Cyperaceae, Rosaceae, Ranunculaceae, Polygonaceae etc. constituting the micro-environment of any one species of *Pedicularis*, each one being specific to their hosts. Further, the roots were intricately entangled

with those of the associated species and formed gregarious patches. The flowers of *Pedicularis* were strikingly delicate and brightly colourful (Fig. 1) and appeared exceptionally prone to perish in the drastic climatic conditions, but species survival perpetuates due to their amazingly adaptive strategies (Garg and Husain, 2003).

The field studies followed by identification in lab and comparative morphological analysis confirmed the existence of 83 species in India with many infra-specific taxa (Husain and Garg, 2007). Out of the existing 83



Fig. 2. Distribution of *Pedicularis* species in India.

Indian species, 81 species find their place on high altitudes of the Himalayan ranges of India and two species, namely *Pedicularis zeylanica* and *P. perrottetii*, are confined to the Nilgiri hills of southern India. The Eastern Himalayas were endowed with an admixture of 44 species, most of them occurring in the Sikkim Himalayas from where they proliferate in all directions, and generally appear as heterogenous assemblage. The number of species gradually decrease westwards and become restricted to 26 species in the Western Himalayas, where they normally form homogenous population patches. There were as many as 11 species which were common to the east and west Himalayan zones (Fig. 3). Most of these were represented in Sikkim (Table 1). A complete list of species that are found in the Sikkim Himalayas, including those endemic to Sikkim, is provided in Table 1. An outstanding amalgam of 49 species (ca. 59%) existed in Sikkim, of which 34 species (41%) were strictly endemic. The maximum species diversity and concentration (49 species) coupled with noticeable endemism (34 species) was found in Sikkim. Due to the strikingly diverse assemblage and rich endemism, the Sikkim Himalayas may be proposed as the 'megadiversity centre' of Indian *Pedicularis*.

ACKNOWLEDGEMENTS

The authors are thankful to the Director, NBRI for providing facilities during the course of this work. The first author (AG) is also grateful to the Director, BSI, Dr. M.




Fig. 3. Statistical representation of Distribution of *Pedicularis* species in India.

Sanjappa and Additional Director, Dr. M.S. Mondal, for facilities and encouragement during publication.

LITERATURE CITED

- Bunge, A. 1846. Ueber *Pedicularis comosa* L. und die mit ihr verwandten Arten. Bull. Phys. Math. Acad. Sci. St. Petersb. **1**: 369-384.
- Garg, A. and T. Husain. 2003. Strategies adopted by the alpine genus *Pedicularis* L. (Scrophulariaceae) to overcome environmental stress. Current Science. **85**: 1413-1414.
- Husain, T. and A. Garg. 2007. Revision of *Pedicularis* L. (Scrophulariaceae) in India. Bishen Singh Mah. Pal Singh print. & publ., Cannaught Place, DehraDun, India. (in press).
- Li, H.-L. 1951. Evolution in the flowers of *Pedicularis*. Evolution **5**: 158-164.
- Linnaeus, C. 1937. *Genera Plantarum*, Leiden. [Ge. Pl.].
- Maximowicz, C. J. 1882. Diagnoses plantarum novarum Asiaticarum. IV. Bull. Acad. Sci. St.-Petersb. **27**: 425-560, pl. 1; in Mel. Biol Acad. Sci. St. Petersb. **11**: 155-350, pl. 1.
- Mill, R. R. 2001. Notes relating to the flora of Bhutan: XLIII. Scrophulariaceae (*Pedicularis*). Edinb. J. Bot. **58**: 57-98.
- Pennell, F. W. 1943. *Pedicularis* In: The Scrophulariaceae of the western Himalayas. The Academy of Natural Sciences of Philadelphia. Monograph **5**: 1-163.
- Prain, D. 1889. Noviciae Indiae. I. Some additional species of *Pedicularis*. J. As. Soc. Bengal **58**: 255-278.
- Prain, D. 1890. The species of *Pedicularis* of the Indian Empire and its Frontiers In: Annals of the Royal Botanic Garden, Calcutta. **1**: 1-196.
- Tsoong, P. C. 1955a. New Himalayan species of *Pedicularis*. Bull. Brit. Mus. (Nat. Hist.) **2**: 1-34.
- Tsoong, P. C. 1955b. Genus *Pedicularis* in Ludlow-Sherriff and Polunion collections. Acta Phytotax. Sin. **3**: 273-344.
- Yang, H.-B., N. H. Holmgren and R. R. Mill. 1998. *Pedicularis* In: Wu, Z.-Y. and P. H. Raven (eds.), Flora of China. **18**: 97-209. Beijing Science Press and St. Louis: Missouri Botanical Garden, China.



錫金喜馬拉雅地區—馬先蒿屬植物在印度之超級多樣性中心

Arti Garg^(1*) and Tariq Husain⁽²⁾

1. Central National Herbarium, Botanical Survey of India, P.O. Indian Botanic Gardens, Howrah 711 103, India. Email: kad_arti396@yahoo.com
2. Taxonomy and Biodiversity Division, National Botanical Research Institute, Rana Pratap Marg, Lucknow – 226 001 (U.P.), India. Email: Hustar_2000@yahoo.co.uk

* 通信作者。

(收稿日期：2009年8月12日；接受日期：2009年11月16日)

摘要：在喜馬拉雅地區馬先蒿屬植物的調查中，顯示本屬植物在錫金喜馬拉雅地區物種相當豐富密集而且具有極高的物種多樣性。印度之 83 種馬先蒿屬植物中，在錫金喜馬拉雅地區就有 49 種(約占 59%)，而且其中 34 種(約占 41%)為本地區之特有種。因此我們可以說錫金喜馬拉雅地區是馬先蒿屬植物在印度之超級多樣性中心。

關鍵詞：馬先蒿屬、玄參科、錫金喜馬拉雅地區、特有的(種)、超級多樣性中心。