

Taiwania*An International Journal of Biodiversity*, established 1947

Editors-in-Chief **Chang-Fu Hsieh, Wen-Yuan Kao**
 Associate Editors **Pei-Chun Liao, Jenn-Che Wang, Roland Kirschner,
 En-Cheng Yang, Yiching Lin, Pei-Jen Lee Shaner,
 Chuan-Kai Ho, Tsung-Hsin Hsieh**
 Managing Editor **Kuo-Hsiung Wang**

Editorial Board

Grecebio Jonathan D. Alejandro ; Philippines	Bai-Ling Lin ; Taiwan
David E. Boufford ; USA	Hiroyoshi Ohashi ; Japan
Su-Hwa Chen ; Taiwan	Arun K. Pandey ; India
Peter L. Chesson ; USA	Chiou-Rong Sheue ; Taiwan
Wen-Liang Chiou ; Taiwan	Shiuh-Feng Shiao ; Taiwan
Chang-Hung Chou ; Taiwan	Kitichate Sridith ; Thailand
Gabor Csorba ; Hungary	Hang Sun ; China
Sauren Das ; India	Iván A. Valdespino ; Panama
Christopher Roy Fraser-Jenkins ; Nepal	Chun-Neng Wang ; Taiwan
Jer-Ming Hu ; Taiwan	Ruijiang Wang ; China
Shih-Tong Jeng ; Taiwan	Jiunn-Tzong Wu ; Taiwan
Yeong-Choy Kam ; Taiwan	Ming-Jou Wu ; Taiwan
Chow-Yang Lee ; USA	Wen-Jer Wu ; Taiwan
Serge Morand ; France	Tsung-Yu Aleck Yang ; Taiwan

Taiwania is an open access (available online at <http://taiwania.ntu.edu.tw>) and peer-reviewed journal issued quarterly, with one volume per year. **The Journal** covers all aspects of biodiversity, with a regional focus on East and Southeast Asia. The content is indexed by major abstracting and indexing services, such as SCIE, Biosis Previews, Zoological Record, Scopus, DOAJ and CAB Abstracts. **Author guidelines** are available on the journal website, and shortened version is also on back cover of this issue. Submitted papers must be original and should not be currently submitted nor published elsewhere. Printed copies can be obtained from the **Taiwania Editorial Office, College of Life Science, National Taiwan University, No. 1, Section 4, Roosevelt Road, Taipei 106, Taiwan.**

ISSN: 0372-333X**<https://taiwania.ntu.edu.tw>**

Cover Photo: *Impatiens bungeilang* Mustaqim is a new species from the northern Gayo Plateau, northern Sumatra. In *Sumatra*, this species alongside *I. calendulina* Grey-Wilson, can be recognized by the whorled leaves and bucciniform lower sepals but the flowers of *I. calendulina* are bright orange-yellow, while in *I. bungeilang*, the flowers are generally red. Another species with red flowers, *I. rubriflora* Grey-Wilson, but this species differs in having spirally arranged leaves.

Acknowledgments: This publication is supported in part by Grants from National Taiwan University and National Science and Technology Council, ROC.