



Supplement

The following supplementary materials are available for this article: Huang, Y.-M., Tseng, M.-H., Chang, Y.-H., Chiou, W.-L., Lee, P.-H. 2025 Factors affecting stipule sprouting of *Angiopteris somae* (Marattiaceae). *Taiwania* 70(2): 361-366. doi: [10.6165/tai.2024.70.361](https://doi.org/10.6165/tai.2024.70.361)

Table S1. Stipule number (SN) and estimated age of plant (Ap), mean stipule width (MW), mean stipule weight (MM), sprouting percentage (SP), mean sprouting number (MSN), and mean sprouting time (MST) of plant stipules in the fern *Angiopteris somae*.

Pl. no.	SN	Ap (years)	MW (mm)	MM (g)	SP (%)	MSN	MST (wks)
1	36	18.3	17.4±3.6 ^{bc}	3.1±1.1 ^{ab}	61.1	0.9±0.8 ^a	15.7±2.8 ^{abc}
2	16	8.1	16.4±5.8 ^{bc}	2.0±0.4 ^b	62.5	0.9±0.9 ^a	13.6±3.0 ^{bc}
3	25	12.7	15.6±2.6 ^c	2.5±0.7 ^b	60.0	0.8±0.8 ^a	17.4±1.8 ^{ab}
4	17	8.6	14.1±3.3 ^{cd}	1.5±0.5 ^c	82.4	1.4±0.8 ^a	16.5±3.5 ^{abc}
5	24	12.2	16.9±3.0 ^{bc}	2.4±0.7 ^b	91.7	1.4±0.7 ^a	14.9±4.0 ^{bc}
6	24	12.2	17.1±2.3 ^{bc}	2.0±0.3 ^b	75.0	1.1±0.8 ^a	16.9±3.2 ^{abc}
7	35	17.8	18.2±4.1 ^a	2.7±1.7 ^b	82.0	1.3±0.8 ^a	15.0±2.5 ^{bc}
8	29	14.7	15.0±3.1 ^{cd}	2.0±1.0 ^b	65.5	1.1±0.9 ^a	16.2±2.7 ^{abc}
9	24	12.2	16.8±3.7 ^{bc}	2.3±1.1 ^b	87.5	1.5±0.7 ^a	14.8±3.5 ^{bc}
10	26	13.2	18.7±4.1 ^{ab}	3.3±1.1 ^{ab}	76.9	1.2±0.8 ^a	15.0±3.5 ^{bc}
11	40	20.3	20.6±3.1 ^a	4.0±1.3 ^a	75.0	1.2±0.8 ^a	15.9±3.2 ^{abc}
12	17	8.6	16.2±2.5 ^{bc}	2.4±0.9 ^b	52.9	0.9±0.9 ^a	18.8±3.2 ^a
13	22	11.2	20.9±2.7 ^a	4.0±1.0 ^a	72.7	1.2±0.9 ^a	13.6±3.3 ^c
14	24	12.2	15.8±2.3 ^c	2.5±0.7 ^b	79.2	1.1±0.7 ^a	15.8±3.7 ^{abc}
15	25	12.7	16.6±2.9 ^{bc}	2.9±0.5 ^b	80.0	1.4±0.8 ^a	14.5±3.2 ^{bc}
16	17	8.6	12.2±5.2 ^d	1.8±0.8 ^b	100.0	1.6±0.5 ^a	14.7±2.8 ^{bc}
17	30	15.2	15.9±2.7 ^c	2.3±0.9 ^b	53.3	0.9±0.9 ^a	17.1±3.5 ^{ab}
18	34	17.3	19.2±5.8 ^b	3.9±1.8 ^a	67.6	1.2±0.9 ^a	14.9±3.8 ^{bc}
19	26	13.2	15.4±3.2 ^{cd}	2.2±0.4 ^b	84.6	1.4±0.8 ^a	14.2±4.2 ^{bc}
20	21	10.7	13.0±2.2 ^d	1.5±0.6 ^{bc}	71.4	1.0±0.8 ^a	13.4±3.6 ^c