

Taiwania*An International Journal of Biodiversity*, established 1947

Editors-in-Chief **Chang-Fu Hsieh, Wen-Yuan Kao**
 Associate Editors **Kuo-Jung Chao, Tsung-Hsin Hsieh, Roland Kirschner,
 Chia-Ying Ko, Pei-Chun Liao, Pei-Jen Lee Shaner,
 Jenn-Che Wang, En-Cheng Yang**
 Managing Editor **Kuo-Hsiung Wang**

Editorial Board

Grecebio Jonathan D. Alejandro ; Philippines	Yiching Lin ; Taiwan
David E. Boufford ; USA	Serge Morand , France
Peter L. Chesson ; USA	Hiroyoshi Ohashi ; Japan
Wen-Liang Chiou ; Taiwan	Arun K. Pandey ; India
Chang-Hung Chou ; Taiwan	Chiou-Rong Sheue ; Taiwan
Gabor Csorba ; Hungary	Shiuh-Feng Shiao ; Taiwan
Sauren Das ; India	Kitichate Sridith ; Thailand
Christopher Roy Fraser-Jenkins ; Nepal	Hang Sun ; China
Chuan-Kai Ho ; Taiwan	Iván A. Valdespino ; Panama
Jer-Ming Hu ; Taiwan	Chun-Neng Wang ; Taiwan
Shih-Tong Jeng ; Taiwan	Ruijiang Wang ; China
Yeong-Choy Kam ; Taiwan	Jiunn-Tzong Wu ; Taiwan
Chow-Yang Lee ; USA	Ming-Jou Wu ; Taiwan
Bai-Ling Lin ; Taiwan	Tsung-Yu Aleck Yang ; Taiwan

Taiwania is an open access (available online at <http://taiwania.ntu.edu.tw>) and peer-reviewed journal issued quarterly, with one volume per year. **The Journal** covers all aspects of biodiversity, with a regional focus on East and Southeast Asia. The content is indexed by major abstracting and indexing services, such as SCIE, Biosis Previews, Zoological Record, Scopus, EBSCO, ProQuest and Ingenta Abstracts. **Author guidelines** are available on the journal website, and shortened version is also on back cover of this issue. Submitted papers must be original and should not be currently submitted nor published elsewhere. Printed copies can be obtained from the **Taiwania Editorial Office, College of Life Science, National Taiwan University, No. 1, Section 4, Roosevelt Road, Taipei 106, Taiwan.**

ISSN: 0372-333X**<https://taiwania.ntu.edu.tw>**

Cover Photo: *Agapetes mingyuaniana* is morphologically similar to *A. variegata* but differs in having ridged branchlets, shorter pedicels, longer corollas, stamens and styles, and anthers without spurs. It is unique among Chinese *Agapetes* species in combining ridged branchlets and corollas longer than 5.5 cm (photo by Hong-Bo Ding).

Acknowledgments: This publication is supported in part by Grants from National Taiwan University and National Science and Technology Council, ROC.